


**Local Realism and GHZ states**  
**Masatsugu Sei Suzuki**  
**Department of Physics, SUNY at Binghamton**  
**(Date: November 15, 2015)**

For many years, everyone thought that Bell had basically exhausted the subject by considering all really interesting situations, and two-spin systems provides the most spectacular quantum violations of local realism. It therefore came as a surprise to many when in 1989 Greenberger, Hone, and Zeilinger (GHZ) showed that systems containing more than two correlated particles may actually exhibit even more dramatic violations of local realism. They involve a sign contradiction (100 % violation) for perfect correlations, while the CHSH inequalities are violated about 40 % (Tsirelson bound = 2 for the CHSH inequality) and deal with situations where the results of measurements are not completely correlated. (F. Laloë, Do we really understand Quantum Mechanics?, Cambridge, 2012).

---

**Anton Zeilinger** (born on 20 May 1945) is an Austrian quantum physicist who in 2008 received the Inaugural Isaac Newton Medal of the Institute of Physics (UK) for "his pioneering conceptual and experimental contributions to the foundations of quantum physics, which have become the cornerstone for the rapidly-evolving field of quantum information". Zeilinger is professor of physics at the University of Vienna and Senior Scientist at the Institute for Quantum Optics and Quantum Information IQOQI at the Austrian Academy of Sciences. Most of his research concerns the fundamental aspects and applications of quantum entanglement.



[http://en.wikipedia.org/wiki/Anton\\_Zeilinger](http://en.wikipedia.org/wiki/Anton_Zeilinger)

**1. Element of reality**

We consider the decay of a simple system into a pair of spin 1/2 particles such as

$$\pi^0 \rightarrow e^+ + e^-,$$

where  $e^+$  is a positron and  $e^-$  is an electron. After the decay products have separated and are very far apart, we measure a component of the spin of one of them. This is the entangled state.

Suppose that  $\hat{S}_z$  of the electron is measured by Alice using the SGz device with  $B//z$  and is found to be equal to  $\hbar/2$ . Then Alice can be sure that  $\hat{S}_z$  of positron will turn out equal to  $-\hbar/2$ , if Bob measures it, since the positron and electron form the entangled state.

Next we consider the different situation. Alice measures the eigenvalue of  $\hat{S}_z$  for the electron by using her SGz device with  $B//z$ . She finds that the eigenvalue of  $\hat{S}_z$  for electron is equal to  $\hbar/2$ . Suppose that Bob measure the eigenvalue of  $\hat{S}_x$  for the positron by using his SGx with  $B//x$ , instead of measuring with the SGz device. What is the eigenvalue of  $\hat{S}_x$  for the positron measured by Bob?

According to quantum mechanics, the probability of finding the state  $|+x\rangle$  is the same as that of finding the state  $|-x\rangle$ . The probability is equal to 1/2 for each case, since

$$P = |\langle +x | -z \rangle|^2 = |\langle -x | -z \rangle|^2 = \frac{1}{2}$$

with

$$|+x\rangle = \frac{1}{\sqrt{2}}[|+z\rangle + |-z\rangle], \quad |-x\rangle = \frac{1}{\sqrt{2}}[|+z\rangle - |-z\rangle]$$

We note that the spin operators  $\hat{S}_z$  and  $\hat{S}_x$  for the positron are not commutable;  $[\hat{S}_z, \hat{S}_x] \neq 0$ . Thus the eigenvalue of  $\hat{S}_x$  cannot be determined definitely, even if the eigenvalue of  $\hat{S}_z$  for the positron can be determined uniquely as  $-\hbar/2$  because of the entangled state.

In the element of reality as defined by EPR theory (local theory), it is assume that all the spin operators are commutable. So all three components of the spin for the positron can be predictable with certainty, if we measure the corresponding spin component of the positron. This claim, however, is incompatible with quantum mechanics, which asserts that at most one spin component of each particle may be definite.

## 2. Local realism and quantum mechanics

We consider two spin 1/2 particles, far apart from each other, in a singlet state. The Bell's state (singlet, spin zero) is given by

$$|\Phi^-\rangle_{12} = \frac{1}{\sqrt{2}}(|+z\rangle_1|-z\rangle_2 - |-z\rangle_1|+z\rangle_2) = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ 1 \\ -1 \\ 0 \end{pmatrix}$$

We know that measurements of  $\hat{\sigma}_{1x}$  and  $\hat{\sigma}_{2x}$ , if performed, shall yield opposite values, that we denote by  $m_{1x}$  and  $m_{2x}$ , respectively. We note that the quantum mechanics asserts that the singlet state (Bell's state) satisfies

$$(\hat{\sigma}_{1x} \otimes \hat{1}_2 + \hat{1}_1 \otimes \hat{\sigma}_{2x})|\Phi^{(-)}\rangle_{12} = 0$$

or

$$m_{1x} = -m_{2x}$$

where  $m_{1x}$  and  $m_{2x}$  are either 1 or -1. Likewise, measurements of  $\hat{\sigma}_{1y}$  and  $\hat{\sigma}_{2y}$ , if performed, shall yield opposite values, that we denote by  $m_{1y}$  and  $m_{2y}$ , respectively. We note that the quantum mechanics asserts that the singlet state (Bell's state) satisfies

$$(\hat{\sigma}_{1y} \otimes \hat{1}_2 + \hat{1}_1 \otimes \hat{\sigma}_{2y})|\Phi^{(-)}\rangle_{12} = 0$$

or

$$m_{1y} = -m_{2y}.$$

where  $m_{1y}$  and  $m_{2y}$  are either 1 or -1. Furthermore, since  $\hat{\sigma}_{1x}$  and  $\hat{\sigma}_{2y}$  commute, and both correspond to elements of reality, their product  $\hat{\sigma}_{1x}\hat{\sigma}_{2y}$  also corresponds to an element of reality. The numerical value assigned to the product  $\hat{\sigma}_{1x}\hat{\sigma}_{2y}$  is the product of the individual numerical values,  $m_{1x}m_{2y}$ . Likewise, the numerical value assigned to the product  $\hat{\sigma}_{1y}\hat{\sigma}_{2x}$  is the product of the individual numerical values,  $m_{1y}m_{2x}$ . These two products must be equal, since

$$m_{1x}m_{2y} = (-m_{2x})(-m_{2y}) = m_{2x}m_{2y}$$

We note that the quantum mechanics asserts that the singlet state (Bell's state) satisfies

$$(\hat{\sigma}_{1x} \otimes \hat{\sigma}_{2y} + \hat{\sigma}_{1y} \otimes \hat{\sigma}_{2x}) \left| \Phi^{(-)} \right\rangle_{12} = 0$$

The proof of this equation will be given later. From this equation, we can predict with certainty that if we measure  $(\hat{\sigma}_{1x} \otimes \hat{\sigma}_{2y} + \hat{\sigma}_{1y} \otimes \hat{\sigma}_{2x})$ , we have

$$m_{1x}m_{2y} + m_{1y}m_{2x} = 0$$

where each operator corresponds to an EPR element of reality.

We note that this equation is totally inconsistent with the equation derived above based on EPR element of reality;  $m_{1x}m_{2y} = m_{1y}m_{2x}$

We note that

$$(\hat{\sigma}_{1x} \otimes \hat{1}_2 + \hat{1}_1 \otimes \hat{\sigma}_{2x}) = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \end{pmatrix}, \quad (\hat{\sigma}_{1y} \otimes \hat{1}_2 + \hat{1}_1 \otimes \hat{\sigma}_{2y}) = \begin{pmatrix} 0 & 0 & -2i & 0 \\ 0 & 0 & 0 & -2i \\ 2i & 0 & 0 & 0 \\ 0 & 2i & 0 & 0 \end{pmatrix}$$

$$(\hat{\sigma}_{1x} \otimes \hat{\sigma}_{2y} + \hat{\sigma}_{1y} \otimes \hat{\sigma}_{2x}) = \begin{pmatrix} 0 & 0 & 0 & -2i \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 2i & 0 & 0 & 0 \end{pmatrix}$$

### 3. Mathematica

Here, using the Mathematica, we show that the singlet state (Bell's state) satisfies the following relations

$$(\hat{\sigma}_{1x} \otimes \hat{1}_2 + \hat{1}_1 \otimes \hat{\sigma}_{2x}) \left| \Phi^{(-)} \right\rangle_{12} = 0$$

$$(\hat{\sigma}_{1y} \otimes \hat{1}_2 + \hat{1}_1 \otimes \hat{\sigma}_{2y}) \left| \Phi^{(-)} \right\rangle_{12} = 0$$

$$(\hat{\sigma}_{1x} \otimes \hat{\sigma}_{2y} + \hat{\sigma}_{1y} \otimes \hat{\sigma}_{2x}) \left| \Phi^{(-)} \right\rangle_{12} = 0$$

$$(\hat{\sigma}_{1y} \otimes \hat{\sigma}_{2z} + \hat{\sigma}_{1z} \otimes \hat{\sigma}_{2y}) \left| \Phi^{(-)} \right\rangle_{12} = 0$$

$$(\hat{\sigma}_{1z} \otimes \hat{\sigma}_{2x} + \hat{\sigma}_{1x} \otimes \hat{\sigma}_{2z}) \left| \Phi^{(-)} \right\rangle_{12} = 0$$

and

$$(\hat{\sigma}_{1x} \otimes \hat{\sigma}_{2x})|\Phi^{(-)}\rangle_{12} = -|\Phi^{(-)}\rangle_{12}$$

$$(\hat{\sigma}_{1y} \otimes \hat{\sigma}_{2y})|\Phi^{(-)}\rangle_{12} = -|\Phi^{(-)}\rangle_{12}$$

$$(\hat{\sigma}_{z1} \otimes \hat{\sigma}_{z2})|\Phi^{(-)}\rangle_{12} = -|\Phi^{(-)}\rangle_{12}$$

**((Mathematica))**

```
Clear["Global`*"];
exp_ * :=
  exp /. {Complex[re_, im_] => Complex[re, -im]};
psi1 =  $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ ;
psi2 =  $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ ;
sigma_x = PauliMatrix[1]; sigma_y = PauliMatrix[2];
sigma_z = PauliMatrix[3];
chi =
   $\frac{1}{\sqrt{2}}$ 
  (KroneckerProduct[psi1, psi2] -
   KroneckerProduct[psi2, psi1]);

A12 =
  (KroneckerProduct[sigma_x, sigma_y] +
   KroneckerProduct[sigma_y, sigma_x]);
A12.chi
{{0}, {0}, {0}, {0}}
```

**A23 =**  
 (KroneckerProduct[ $\sigma_y$ ,  $\sigma_z$ ] +  
 KroneckerProduct[ $\sigma_z$ ,  $\sigma_y$ ]);

**A23. $\chi$**

{{0}, {0}, {0}, {0}}

**A31 =**  
 (KroneckerProduct[ $\sigma_x$ ,  $\sigma_z$ ] +  
 KroneckerProduct[ $\sigma_z$ ,  $\sigma_x$ ]);

**A31. $\chi$**

{{0}, {0}, {0}, {0}}

(KroneckerProduct[ $\sigma_x$ ,  $\sigma_x$ ]). $\chi$  +  $\chi$

{{0}, {0}, {0}, {0}}

(KroneckerProduct[ $\sigma_y$ ,  $\sigma_y$ ]). $\chi$  +  $\chi$


{{0}, {0}, {0}, {0}}

(KroneckerProduct[ $\sigma_z$ ,  $\sigma_z$ ]). $\chi$  +  $\chi$

{{0}, {0}, {0}, {0}}

---

**4. GHZ state  $|\psi_{GHZ}^-\rangle$  for the spin 1/2 systems**



**Fig.** A source (S) of particle triples produces three identical particles which then move towards three equidistant magnetic orientation detectors. Alice, Bob and Chris set up detectors to measure the magnetization along the direction of the particle's motion (the z-direction) or along two other mutually perpendicular directions (x and y-directions).

In 1989, a striking extension of Bell's theorem to the case of three particles was taken by Greenberger, Horne, and Zeilinger. In contrast to Bell's theorem, which concerns statistical averages, this so-called GHZ theorem shows that a conflict between quantum mechanics and local realism can be obtained with a single measurements. GHZ consider three observers, Alice (1), Bob (2), and Chris (3). The GHZ experiments are a class of physics experiments that may be used to generate starkly contrasting predictions from local hidden variable theory and quantum mechanics, and permits immediate comparison with actual experimental results.

Using the Mathematica, here we show that the singlet state (GHZ state) are the eigenkets of the following operators

$$\hat{A}_{xyy} = \hat{\sigma}_{x1} \otimes \hat{\sigma}_{y2} \otimes \hat{\sigma}_{y3},$$

$$\hat{A}_{yyx} = \hat{\sigma}_{y1} \otimes \hat{\sigma}_{x2} \otimes \hat{\sigma}_{y3},$$

$$\hat{A}_{yxy} = \hat{\sigma}_{y1} \otimes \hat{\sigma}_{y2} \otimes \hat{\sigma}_{x3},$$

$$\hat{A}_{xxx} = \hat{\sigma}_{x1} \otimes \hat{\sigma}_{x2} \otimes \hat{\sigma}_{x3}.$$

First we consider the GHZ state  $|\psi_{GHZ}^-\rangle$  given by

$$|\psi_{GHZ}^-\rangle = \frac{1}{\sqrt{2}} (|+z\rangle_1 | +z\rangle_2 | +z\rangle_3 - |-z\rangle_1 |-z\rangle_2 |-z\rangle_3) = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ -1 \end{pmatrix}$$

It is seen that  $|\psi_{GHS}^-\rangle$  is an eigenstate of several operators with the eigenvalue +1,

$$\hat{A}_{xyy} |\psi_{GHS}^-\rangle = |\psi_{GHS}^-\rangle, \quad (1a)$$

$$\hat{A}_{yyx} |\psi_{GHS}^-\rangle = |\psi_{GHS}^-\rangle, \quad (1b)$$

$$\hat{A}_{yxy} |\psi_{GHS}^-\rangle = |\psi_{GHS}^-\rangle, \quad (1c)$$

$$(1d)$$

Then we obtain

$$\hat{A}_{xyy} \hat{A}_{yyx} \hat{A}_{yxy} |\psi_{GHZ}^-\rangle = |\psi_{GHZ}^-\rangle,$$

We can also show that  $|\psi_{GHS}^-\rangle$  is an eigenstate of the operator  $\hat{A}_{xxx}$  with the eigenvalue -1,



$$\hat{A}_{xxx}|\psi_{GHS}^-\rangle = -|\psi_{GHS}^-\rangle$$

Once three particles are sufficiently far apart, each spin of them possesses its own physical characteristics. We use  $A_x$  to denote the result of measuring the  $x$  component of the spin of particle 1 by Alice,  $B_y$  the result of measuring the  $y$  component of the spin of particle 2 by Bob, and  $C_y$  the result of measuring the  $y$  component of the spin of particle 3 by Chris, and so on, with  $A_x = \pm 1 \dots, C_y = \pm 1$ . When the  $x$  component is measured in connection with two measurements of the  $y$  component, we see that the product is +1:

$$A_x B_y C_y = +1,$$

Similarly, we have

$$A_y B_x C_y = +1, \quad A_y B_y C_x = +1$$

However, when the particles are in flight, two of the three experimentalists can decide to modify the direction of their analyzer axes, orienting them in the  $x$  axis direction. Then the product of the three spin components will be -1:

$$A_x B_x C_x = -1 \tag{2}$$

However, we note that

$$A_x B_x C_x = (A_x B_y C_y)(A_y B_x C_y)(A_y B_y C_x) = 1 \tag{3}$$

because  $A_y^2 = B_y^2 = C_y^2 = 1$ . Thus Eqs.(2) and (3) are incompatible.

Local realism would mean that  $\hat{\sigma}_{x1}$  has a physical reality in the EPR sense, since it can be measured without disturbing  $\hat{\sigma}_{y2}$  and  $\hat{\sigma}_{y3}$ ,

$$A_x = B_y C_y.$$

However, it is also possible to obtain  $A_x$  by measuring  $\hat{\sigma}_{x2}$  and  $\hat{\sigma}_{x3}$ :

$$A_x = -B_x C_x$$

Local realism implies that it is the same  $A_x$ , but this is not the case in quantum mechanics. The value of  $A_x$  is contextual. It depends on physical properties incompatible with each other which are measured simultaneously.

**((Mathematica))**

```

Clear["Global`*"];  $\psi_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}; \psi_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix};$ 
 $\sigma_x = \text{PauliMatrix}[1]; \sigma_y = \text{PauliMatrix}[2];$ 
 $\sigma_z = \text{PauliMatrix}[3];$ 
 $\chi =$ 
 $\frac{1}{\sqrt{2}} (\text{KroneckerProduct}[\psi_1, \psi_1, \psi_1] -$ 
 $\text{KroneckerProduct}[\psi_2, \psi_2, \psi_2]);$ 

 $A_{xyy} = \text{KroneckerProduct}[\sigma_x, \sigma_y, \sigma_y]; A_{xyy} \cdot \chi - \chi$ 
{{0}, {0}, {0}, {0}, {0}, {0}, {0}, {0}}

 $A_{yxy} = \text{KroneckerProduct}[\sigma_y, \sigma_x, \sigma_y]; A_{yxy} \cdot \chi - \chi$ 
{{0}, {0}, {0}, {0}, {0}, {0}, {0}, {0}}

 $A_{yyx} = \text{KroneckerProduct}[\sigma_y, \sigma_y, \sigma_x]; A_{yyx} \cdot \chi - \chi$ 
{{0}, {0}, {0}, {0}, {0}, {0}, {0}, {0}}

 $A_{xxx} = \text{KroneckerProduct}[\sigma_x, \sigma_x, \sigma_x]; A_{xxx} \cdot \chi + \chi$ 
{{0}, {0}, {0}, {0}, {0}, {0}, {0}, {0}}

 $A_{123} = A_{xyy} \cdot A_{yxy} \cdot A_{yyx}; A_{123} \cdot \chi + A_{xxx} \cdot \chi$ 
{{0}, {0}, {0}, {0}, {0}, {0}, {0}, {0}}

```

---

## 5. $|\psi_{GHZ}^+\rangle$ state for spin 1/2 systems

We consider the GHZ state  $|\psi_{GHZ}^+\rangle$  defined by

$$|\psi_{GHZ}^+\rangle = \frac{1}{\sqrt{2}}(|+z\rangle_1|+z\rangle_2|+z\rangle_3 + |-z\rangle_1|-z\rangle_2|-z\rangle_3) = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}$$

It is seen that  $|\psi_{GHS}^+\rangle$  is an eigenstate of several operators with the eigenvalue -1,

$$\hat{A}_{xyy}|\psi_{GHS}^+\rangle = -|\psi_{GHS}^+\rangle, \quad (1a)$$

$$\hat{A}_{yyx}|\psi_{GHS}^+\rangle = -|\psi_{GHS}^+\rangle, \quad (1b)$$

$$\hat{A}_{xyx}|\psi_{GHS}^+\rangle = -|\psi_{GHS}^+\rangle. \quad (1c)$$

Then we obtain

$$\hat{A}_{xyy}\hat{A}_{yyx}\hat{A}_{xyx}|\psi_{GHS}^+\rangle = -\hat{A}_{xyy}\hat{A}_{yyx}|\psi_{GHS}^+\rangle = \hat{A}_{xyy}|\psi_{GHS}^+\rangle = -|\psi_{GHS}^+\rangle, \quad (2)$$

We can also show that  $|\psi_{GHS}^+\rangle$  is an eigenstate of  $\hat{A}_{xxx}$  with the eigenvalue +1,

$$\hat{A}_{xxx}|\psi_{GHS}^+\rangle = |\psi_{GHS}^+\rangle.$$

Once the three particles are sufficiently far apart, each spin of them possesses its own physical characteristics. We use  $A_x$  to denote the result of measuring the  $x$  component of the spin of particle 1 by Alice, ...,  $C_y$  the result of measuring the  $y$  component of the spin of particle 3 by Charlotte, and so on, with  $A_x = \pm 1$  ...,  $C_y = \pm 1$ . When the  $x$  component is measured in connection with two measurements of the  $y$  component, we see that the products is +1:

$$A_x B_y C_y = -1, \quad A_y B_x C_y = -1, \quad A_y B_y C_x = -1$$

However, when the particles are in flight, two of the three experimentalists can decide to modify the direction of their analyzer axes, orienting them in the  $x$  axis direction. Then the product of the three spin components will be -1:

$$A_x B_x C_x = 1 \quad (2)$$

However, we note that

$$A_x B_x C_x = (A_x B_y C_y)(A_y B_x C_y)(A_y B_y C_x) = -1 \quad (3)$$

because  $A_y^2 = B_y^2 = C_y^2 = 1$ . Thus Eqs.(2) and (3) are incompatible. Local realism would mean that  $\hat{\sigma}_{x1}$  has a physical reality in the EPR sense, since it can be measured without disturbing  $\hat{\sigma}_{y2}$  and  $\hat{\sigma}_{y3}$ ,

$$A_x = -B_y C_y .$$

However, it is also possible to obtain  $A_x$  by measuring  $\hat{\sigma}_{x2}$  and  $\hat{\sigma}_{x3}$ :

$$A_x = B_x C_x .$$

Local realism implies that it is the same  $A_x$ , but this is not the case in quantum mechanics. The value of  $A_x$  is contextual. It depends on physical properties incompatible with each other which are measured simultaneously

**((Mathematica))**

```

Clear["Global`*"];  $\psi_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ ;  $\psi_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$ ;  $\sigma_x = \text{PauliMatrix}[1]$ ;
 $\sigma_y = \text{PauliMatrix}[2]$ ;  $\sigma_z = \text{PauliMatrix}[3]$ ;
 $\chi = \frac{1}{\sqrt{2}} (\text{KroneckerProduct}[\psi_1, \psi_1, \psi_1] + \text{KroneckerProduct}[\psi_2, \psi_2, \psi_2])$ ;

 $A_{xyy} = \text{KroneckerProduct}[\sigma_x, \sigma_y, \sigma_y]$ ;  $A_{xyy} \cdot \chi + \chi$ 
{{0}, {0}, {0}, {0}, {0}, {0}, {0}, {0}}

 $A_{yxy} = \text{KroneckerProduct}[\sigma_y, \sigma_x, \sigma_y]$ ;  $A_{yxy} \cdot \chi + \chi$ 
{{0}, {0}, {0}, {0}, {0}, {0}, {0}, {0}}

 $A_{yyx} = \text{KroneckerProduct}[\sigma_y, \sigma_y, \sigma_x]$ ;  $A_{yyx} \cdot \chi + \chi$ 
{{0}, {0}, {0}, {0}, {0}, {0}, {0}, {0}}

 $A_{xxx} = \text{KroneckerProduct}[\sigma_x, \sigma_x, \sigma_x]$ ;  $A_{xxx} \cdot \chi - \chi$ 
{{0}, {0}, {0}, {0}, {0}, {0}, {0}, {0}}

 $A_{123} = A_{xyy} \cdot A_{yxy} \cdot A_{yyx}$ ;  $A_{123} \cdot \chi + A_{xxx} \cdot \chi$ 
{{0}, {0}, {0}, {0}, {0}, {0}, {0}, {0}}

 $A_{xyy} \cdot \chi + \chi$ 
{{0}, {0}, {0}, {0}, {0}, {0}, {0}, {0}}

 $A_{yxy} \cdot \chi + \chi$ 
{{0}, {0}, {0}, {0}, {0}, {0}, {0}, {0}}

 $A_{yyx} \cdot \chi + \chi$ 
{{0}, {0}, {0}, {0}, {0}, {0}, {0}, {0}}

 $A_{xxx} \cdot \chi - \chi$ 
{{0}, {0}, {0}, {0}, {0}, {0}, {0}, {0}}

```

---


## 6. GHZ state for photon systems

Let us denote  $|H\rangle$  by matrix  $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$  and  $|V\rangle$  by matrix  $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ ; they are thus the two eigenstates of the Pauli operator,  $\hat{\sigma}_z$ , correspondingly with the eigenvalues +1 and -1. We can also easily verify that  $|H\rangle$  and  $|V\rangle$  or  $|R\rangle$  and  $|L\rangle$  are two eigenstates for the Pauli operator  $\hat{\sigma}_x$  or  $\hat{\sigma}_y$  with the values +1 and -1, respectively. For convenience we will refer to a measurement of the  $H'/V'$  linear polarization as an x measurement and of the  $L/R$  circular polarization as a y measurement.

**((Note)) Pauli matrices for photon polarization**

For convenience we use the Pauli matrices

$$\hat{\sigma}_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \hat{\sigma}_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \hat{\sigma}_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$



(a) The Pauli operator  $\hat{\sigma}_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ :

$|H'\rangle$  and  $|V'\rangle$  are the eigenkets of  $\hat{\sigma}_x$  with the eigenvalues +1, and -1, respectively.

$$\hat{\sigma}_x |H'\rangle = |H'\rangle, \quad \hat{\sigma}_x |V'\rangle = -|V'\rangle$$

(b) The Pauli operator  $\hat{\sigma}_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$ :

$|R\rangle$  and  $|L\rangle$  are the eigenkets of  $\hat{\sigma}_y$  with the eigenvalues +1, and -1, respectively.

$$\hat{\sigma}_y |R\rangle = |R\rangle, \quad \hat{\sigma}_y |L\rangle = -|L\rangle$$

(c) The Pauli operator  $\hat{\sigma}_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$

$|x\rangle$  and  $|y\rangle$  are the eigenkets of  $\hat{\sigma}_z$  with the eigenvalues +1, and -1, respectively.

((**Comparison**)) photon and spin states

$$|45^\circ\rangle = |H'\rangle = \frac{1}{\sqrt{2}}(|x\rangle + |y\rangle)$$

$$|+x\rangle = \frac{1}{\sqrt{2}}(|+z\rangle + |-z\rangle)$$

$$|-45^\circ\rangle = |V'\rangle = \frac{1}{\sqrt{2}}(|x\rangle - |y\rangle)$$

$$|-x\rangle = \frac{1}{\sqrt{2}}(|+z\rangle - |-z\rangle)$$

$$|R\rangle = \frac{1}{\sqrt{2}}(|x\rangle + i|y\rangle)$$

$$|+y\rangle = \frac{1}{\sqrt{2}}(|+z\rangle + i|-z\rangle)$$

$$|L\rangle = \frac{1}{\sqrt{2}}(|x\rangle - i|y\rangle)$$

$$|-y\rangle = \frac{1}{\sqrt{2}}(|+z\rangle - i|-z\rangle)$$

## 7. Expressions of the three photon state

(a) Basis  $\{|H'\rangle, |V'\rangle\}$

$$|\psi^{(+)}\rangle = \frac{1}{\sqrt{2}}(|x\rangle_1 |x\rangle_2 |y\rangle_3 + |y\rangle_1 |y\rangle_2 |x\rangle_3)$$

where  $|x\rangle$  and  $|y\rangle$  denote horizontal and vertical polarizations, respectively. This state indicates that the three photons are in a quantum superposition of the state  $|x\rangle_1|x\rangle_2|y\rangle_3$  and  $|y\rangle_1|y\rangle_2|x\rangle_3$ . Note that

$$|x\rangle = \frac{1}{\sqrt{2}}[|H'\rangle + |V'\rangle], \quad |y\rangle = \frac{1}{\sqrt{2}}[|H'\rangle - |V'\rangle]$$

Then we have the form

$$|\psi^{(+)}\rangle = \frac{1}{2}(|H'\rangle_1|H'\rangle_2|H'\rangle_3 - |H'\rangle_1|V'\rangle_2|V'\rangle_3 - |V'\rangle_1|H'\rangle_2|V'\rangle_3 + |V'\rangle_1|V'\rangle_2|H'\rangle_3)$$

((Mathematica))



```

Clear["Global`*"]; x1 =  $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ ; y1 =  $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ ;

 $\chi =$ 
 $\frac{1}{\sqrt{2}}$  (KroneckerProduct[x1, x1, y1] +
    KroneckerProduct[y1, y1, x1]);

H1 =  $\frac{1}{\sqrt{2}}$   $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$ ; V1 =  $\frac{1}{\sqrt{2}}$   $\begin{pmatrix} 1 \\ -1 \end{pmatrix}$ ;

HHH1 = KroneckerProduct[H1, H1, H1] // Simplify;
HHV1 = KroneckerProduct[H1, H1, V1] // Simplify;
HVH1 = KroneckerProduct[H1, V1, H1] // Simplify;
HVV1 = KroneckerProduct[H1, V1, V1] // Simplify;
VHH1 = KroneckerProduct[V1, H1, H1] // Simplify;
VHV1 = KroneckerProduct[V1, H1, V1] // Simplify;
VVH1 = KroneckerProduct[V1, V1, H1] // Simplify;
VVV1 = KroneckerProduct[V1, V1, V1] // Simplify;

f1 = a1 HHH1 + a2 HHV1 + a3 HVH1 + a4 HVV1 +
    a5 VHH1 + a6 VHV1 + a7 VVH1 + a8 VVV1 // Simplify;

eq1 = Solve[f1 ==  $\chi$ , {a1, a2, a3, a4, a5, a6, a7, a8}];

rule1 = {b1 → HHH, b2 → HHV, b3 → HVH, b4 → HVV,
    b5 → VHH, b6 → VHV, b7 → VVH, b8 → VVV};
P1 = a1 b1 + a2 b2 + a3 b3 + a4 b4 + a5 b5 +
    a6 b6 + a7 b7 + a8 b8;
P1 /. rule1 /. eq1[[1]]

 $\frac{HHH}{2} - \frac{HVV}{2} - \frac{VHV}{2} + \frac{VVH}{2}$ 

```

---

(b) Basis  $\{|H'\rangle, |V'\rangle\}$

$$|\psi^{(+)}\rangle = \frac{1}{\sqrt{2}} (|x\rangle_1 |x\rangle_2 |x\rangle_3 + |y\rangle_1 |y\rangle_2 |y\rangle_3)$$

where  $|x\rangle$  and  $|y\rangle$  denote horizontal and vertical polarizations, respectively. This state indicates that the three photons are in a quantum superposition of the state  $|x\rangle_1|x\rangle_2|x\rangle_3$  and  $|y\rangle_1|y\rangle_2|y\rangle_3$ . Using the basis of  $\{|H'\rangle, |V'\rangle\}$ ,  $|\psi^{(+)}\rangle$  can be rewritten as

$$|\psi^{(+)}\rangle = \frac{1}{2}(|H'\rangle_1|H'\rangle_2|H'\rangle_3 + |H'\rangle_1|V'\rangle_2|V'\rangle_3 + |V'\rangle_1|H'\rangle_2|V'\rangle_3 + |V'\rangle_1|V'\rangle_2|H'\rangle_3)$$

((Mathematica))

```
Clear["Global`*"]; x1 =  $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ ; y1 =  $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ ;
χ =
 $\frac{1}{\sqrt{2}}$  (KroneckerProduct[x1, x1, x1] +
KroneckerProduct[y1, y1, y1]);

H1 =  $\frac{1}{\sqrt{2}}$   $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$ ; V1 =  $\frac{1}{\sqrt{2}}$   $\begin{pmatrix} 1 \\ -1 \end{pmatrix}$ ;
HHH1 = KroneckerProduct[H1, H1, H1] // Simplify;
HHV1 = KroneckerProduct[H1, H1, V1] // Simplify;
HVH1 = KroneckerProduct[H1, V1, H1] // Simplify;
HVV1 = KroneckerProduct[H1, V1, V1] // Simplify;
VHH1 = KroneckerProduct[V1, H1, H1] // Simplify;
VHV1 = KroneckerProduct[V1, H1, V1] // Simplify;
VVH1 = KroneckerProduct[V1, V1, H1] // Simplify;
VVV1 = KroneckerProduct[V1, V1, V1] // Simplify;

f1 = a1 HHH1 + a2 HHV1 + a3 HVH1 + a4 HVV1 +
a5 VHH1 + a6 VHV1 + a7 VVH1 + a8 VVV1 // Simplify;

eq1 = Solve[f1 == χ, {a1, a2, a3, a4, a5, a6, a7, a8}];

rule1 = {b1 → HHH, b2 → HHV, b3 → HVH, b4 → HVV,
b5 → VHH, b6 → VHV, b7 → VVH, b8 → VVV};
P1 = a1 b1 + a2 b2 + a3 b3 + a4 b4 + a5 b5 +
a6 b6 + a7 b7 + a8 b8;
P1 /. rule1 /. eq1[[1]]

 $\frac{HHH}{2} + \frac{HVV}{2} + \frac{VHV}{2} + \frac{VVH}{2}$ 
```

(c) Basis  $\{|R\rangle, |L\rangle\}$

$$|\psi^{(-)}\rangle = \frac{1}{\sqrt{2}}(|x\rangle_1|x\rangle_2|x\rangle_3 - |y\rangle_1|y\rangle_2|x\rangle_3)$$

where  $|x\rangle$  and  $|y\rangle$  denote horizontal and vertical polarizations, respectively. This state indicates that the three photons are in a quantum superposition of the state  $|x\rangle_1|x\rangle_2|x\rangle_3$  and  $|y\rangle_1|y\rangle_2|x\rangle_3$ .

Using the basis of  $\{|R\rangle, |L\rangle\}$ ,  $|\psi^{(-)}\rangle$  can be rewritten as

$$|\psi^{(-)}\rangle = \frac{1}{2}(|R\rangle_1|R\rangle_2|R\rangle_3 + |R\rangle_1|R\rangle_2|L\rangle_3 + |L\rangle_1|L\rangle_2|R\rangle_3 + |L\rangle_1|L\rangle_2|L\rangle_3)$$

((**Mathematica**))

```

Clear["Global`*"]; x1 =  $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ ; y1 =  $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ ;

χ =
 $\frac{1}{\sqrt{2}}$  (KroneckerProduct[x1, x1, x1] -
KroneckerProduct[y1, y1, x1]);

R1 =  $\frac{1}{\sqrt{2}}$   $\begin{pmatrix} 1 \\ i \end{pmatrix}$ ; L1 =  $\frac{1}{\sqrt{2}}$   $\begin{pmatrix} 1 \\ -i \end{pmatrix}$ ;

RRR1 = KroneckerProduct[R1, R1, R1] // Simplify;
RRL1 = KroneckerProduct[R1, R1, L1] // Simplify;
RLR1 = KroneckerProduct[R1, L1, R1] // Simplify;
RLL1 = KroneckerProduct[R1, L1, L1] // Simplify;
LRR1 = KroneckerProduct[L1, R1, R1] // Simplify;
LRL1 = KroneckerProduct[L1, R1, L1] // Simplify;
LLR1 = KroneckerProduct[L1, L1, R1] // Simplify;
LLL1 = KroneckerProduct[L1, L1, L1] // Simplify;

f1 = a1 RRR1 + a2 RRL1 + a3 RLR1 + a4 RLL1 +
a5 LRR1 + a6 LRL1 + a7 LLR1 + a8 LLL1 // Simplify;

eq1 = Solve[f1 == χ, {a1, a2, a3, a4, a5, a6, a7, a8}];

rule1 = {b1 → RRR, b2 → RRL, b3 → RLR, b4 → RLL,
b5 → LRR, b6 → LRL, b7 → LLR, b8 → LLL};
P1 = a1 b1 + a2 b2 + a3 b3 + a4 b4 + a5 b5 +
a6 b6 + a7 b7 + a8 b8;
P1 /. rule1 /. eq1[[1]]

 $\frac{LLL}{2} + \frac{LLR}{2} + \frac{RRL}{2} + \frac{RRR}{2}$ 

```

## 8. General case


$$|\psi^{(+)}\rangle = \frac{1}{\sqrt{2}} (|x\rangle_1 |x\rangle_2 |x\rangle_3 + |y\rangle_1 |y\rangle_2 |y\rangle_3)$$

$$|H'\rangle = \cos\theta |x\rangle + \sin\theta |y\rangle,$$

$$|V'\rangle = -\sin\theta|x\rangle + \cos\theta|y\rangle,$$

$$\begin{aligned} |\psi^{(+)}\rangle &= \frac{1}{\sqrt{2}}(\cos^3\theta + \sin^3\theta)|H'\rangle_1|H'\rangle_2|H'\rangle_3 \\ &+ \frac{1}{\sqrt{2}}(\cos^3\theta - \sin^3\theta)|V'\rangle_1|V'\rangle_2|V'\rangle_3 \\ &+ \frac{1}{\sqrt{2}}\sin\theta(\sin\theta - \cos\theta)(|H'\rangle_1|H'\rangle_2|V'\rangle_3 + |H'\rangle_1|V'\rangle_2|H'\rangle_3 + |V'\rangle_1|H'\rangle_2|H'\rangle_3) \\ &+ \frac{1}{\sqrt{2}}\sin\theta(\sin\theta + \cos\theta)(|H'\rangle_1|V'\rangle_2|V'\rangle_3 + |V'\rangle_1|V'\rangle_2|H'\rangle_3 + |V'\rangle_1|H'\rangle_2|V'\rangle_3) \end{aligned}$$

We make a plot of the probabilities for the eight states  $\{|H'\rangle_1|H'\rangle_2|H'\rangle_3, |V'\rangle_1|V'\rangle_2|V'\rangle_3, \dots\}$  as a function of  $\theta$ .



**Fig.** plot of the probabilities for the eight states  $\{|H'\rangle_1|H'\rangle_2|H'\rangle_3, |V'\rangle_1|V'\rangle_2|V'\rangle_3, \dots\}$  as a function of  $\theta$ .

## 9. Reality

For convenience we will refer to a measurement of the  $H'/V'$  linear polarization as an  $x$  measurement and of the  $L/R$  circular polarization as a  $y$  measurement. We have three particles, and we can choose to measure each on arbitrary basis. We designate a chosen set of observation

on these particles by a sequence of symbol  $x$  and  $y$ . If we may choose to measure particles 1, 2, and 3 in only the basis  $x$ . This measurement is denoted by  $xxx$ .

$$\begin{aligned} |\psi^{(+)}\rangle &= \frac{1}{\sqrt{2}}(|x\rangle_1|x\rangle_2|x\rangle_3 + |y\rangle_1|y\rangle_2|y\rangle_3) \\ &= \frac{1}{2}(|H_1'\rangle|H_2'\rangle|H_3'\rangle + |H_1'\rangle|V_2'\rangle|V_3'\rangle + |V_1'\rangle|H_2'\rangle|V_3'\rangle + |V_1'\rangle|V_2'\rangle|H_3'\rangle) \end{aligned}$$

We may also choose to measure particles 1 and 2 in basis  $y$  and particle 3 in basis  $x$ . This measurement is denoted by  $yyx$ . The state  $|\psi\rangle$  expressed in the corresponding basis set becomes

$$\begin{aligned} |\psi^{(+)}\rangle &= \frac{1}{\sqrt{2}}(|y\rangle_1|y\rangle_2|x\rangle_3 + |x\rangle_1|x\rangle_2|y\rangle_3) \\ &= \frac{1}{2}[|R_1'\rangle \otimes |L_2'\rangle \otimes |H_3'\rangle + |L_1'\rangle \otimes |R_2'\rangle \otimes |H_3'\rangle \\ &\quad + |R_1'\rangle \otimes |R_2'\rangle \otimes |V_3'\rangle + |L_1'\rangle \otimes |L_2'\rangle \otimes |V_3'\rangle] \end{aligned}$$

**((Mathematica))**

```

Clear["Global`*"]; x1 =  $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ ; y1 =  $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ ;

H11 =  $\frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix}$ ; v11 =  $\frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ -1 \end{pmatrix}$ ; R1 =  $\frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ i \end{pmatrix}$ ;
L1 =  $\frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ -i \end{pmatrix}$ 

{{ $\frac{1}{\sqrt{2}}$ }, {- $\frac{i}{\sqrt{2}}$ }}

x =
 $\frac{1}{2}$  (KroneckerProduct[R1, L1, H11] +
      KroneckerProduct[L1, R1, H11] +
      KroneckerProduct[R1, R1, V11] +
      KroneckerProduct[L1, L1, V11]);

x // Simplify


{{ $\frac{1}{\sqrt{2}}$ }, {0}, {0}, {0}, {0}, {0}, {0}, { $\frac{1}{\sqrt{2}}$ }}
```

## 10. $yyx$ -, $yxy$ -, and $xyy$ -type measurements

### (a) $xyy$ measurement

The state given by


$$\begin{aligned} |\psi^{xyy(+)}\rangle &= \frac{1}{\sqrt{2}}(|x_1\rangle|y_2\rangle|y_3\rangle + |y_1\rangle|x_2\rangle|x_3\rangle) \\ &= \frac{1}{2}[|H_1'\rangle \otimes |R_2\rangle \otimes |L_3\rangle + |H_1'\rangle \otimes |L_2\rangle \otimes |R_3\rangle \\ &\quad + |V_1'\rangle \otimes |R_2\rangle \otimes |R_3\rangle + |V_1'\rangle \otimes |L_2\rangle \otimes |L_3\rangle] \end{aligned}$$



**Fig.** Fraction of the various outcomes observed in the  $xyy$  measurement (**Pan et.al.**)

### (b) $yxy$ measurement

$$\begin{aligned} |\psi^{yxy(+)}\rangle &= \frac{1}{\sqrt{2}}(|y_1\rangle|x_2\rangle|y_3\rangle + |y_1\rangle|x_2\rangle|y_3\rangle) \\ &= \frac{1}{2}[|L_1\rangle \otimes |H_2'\rangle \otimes |R_3\rangle + |R_1\rangle \otimes |H_2'\rangle \otimes |L_3\rangle \\ &\quad + |R_1\rangle \otimes |V_2'\rangle \otimes |R_3\rangle + |L_1\rangle \otimes |V_2'\rangle \otimes |L_3\rangle] \end{aligned}$$




**Fig.** Fraction of the various outcomes observed in the  $yxy$  measurement (**Pan et.al.**)

**(c)  $yyx$  measurement**

The state for the  $yyx$  measurement is given by

$$\begin{aligned}
 |\psi^{yyx(+)}\rangle &= \frac{1}{\sqrt{2}}(|y_1\rangle|y_2\rangle|x_3\rangle + |x_1\rangle|x_2\rangle|y_3\rangle) \\
 &= \frac{1}{2} [ |R_1\rangle \otimes |L_2\rangle \otimes |H_3'\rangle + |L_1\rangle \otimes |R_2\rangle \otimes |H_3'\rangle \\
 &\quad + |R_1\rangle \otimes |R_2\rangle \otimes |V_3'\rangle + |L_1\rangle \otimes |L_2\rangle \otimes |V_3'\rangle ]
 \end{aligned}$$



**Fig.** Fraction of the various outcomes observed in the  $yyx$  measurement (**Pan et.al.**)

**11. The eigenstates of  $\hat{\sigma}_{1x} \otimes \hat{\sigma}_{2x} \otimes \hat{\sigma}_{3x}$  :  $xxx$ -type measurements**

We consider the eigenstates of  $\hat{\sigma}_{1x} \otimes \hat{\sigma}_{2x} \otimes \hat{\sigma}_{3x}$ . In quantum mechanics, we get



$$\hat{\sigma}_{1x} \otimes \hat{\sigma}_{2x} \otimes \hat{\sigma}_{3x} |V_1'\rangle \otimes |V_2'\rangle \otimes |V_3'\rangle = -|V_1'\rangle \otimes |V_2'\rangle \otimes |V_3'\rangle$$

$$\hat{\sigma}_{1x} \otimes \hat{\sigma}_{2x} \otimes \hat{\sigma}_{3x} |H_1'\rangle \otimes |H_2'\rangle \otimes |V_3'\rangle = -|H_1'\rangle \otimes |H_2'\rangle \otimes |V_3'\rangle$$


$$\hat{\sigma}_{1x} \otimes \hat{\sigma}_{2x} \otimes \hat{\sigma}_{3x} |H_1'\rangle \otimes |V_2'\rangle \otimes |H_3'\rangle = -|H_1'\rangle \otimes |V_2'\rangle \otimes |H_3'\rangle$$

$$\hat{\sigma}_{1x} \otimes \hat{\sigma}_{2x} \otimes \hat{\sigma}_{3x} |V_1'\rangle \otimes |H_2'\rangle \otimes |H_3'\rangle = -|V_1'\rangle \otimes |H_2'\rangle \otimes |H_3'\rangle$$

Then we find that there are four eigenstates of  $\hat{\sigma}_{1x} \otimes \hat{\sigma}_{2x} \otimes \hat{\sigma}_{3x}$

$$|V_1'\rangle \otimes |V_2'\rangle \otimes |V_3'\rangle, |H_1'\rangle \otimes |H_2'\rangle \otimes |V_3'\rangle, \\ |H_1'\rangle \otimes |V_2'\rangle \otimes |H_3'\rangle, \text{ and } |V_1'\rangle \otimes |H_2'\rangle \otimes |H_3'\rangle$$

which have the same eigenvalue (-1).



**Fig.** Prediction from the **local hidden theory**. There are four states denoted by  $|V_1'\rangle \otimes |V_2'\rangle \otimes |V_3'\rangle$ ,  $|H_1'\rangle \otimes |H_2'\rangle \otimes |V_3'\rangle$ ,  $|H_1'\rangle \otimes |V_2'\rangle \otimes |H_3'\rangle$ , and  $|V_1'\rangle \otimes |H_2'\rangle \otimes |H_3'\rangle$ . **(Pan et.al.)**

Thus any superposition of these states has the eigenvalue (-1). However, there is only one GHZ state, among these states, which is defined by

$$\begin{aligned} |\Psi_{GHZ}^{xxx(-)}\rangle &= \frac{1}{\sqrt{2}}[|x_1\rangle|x_2\rangle|x_3\rangle - |y_1\rangle|y_2\rangle|y_3\rangle] \\ &= \frac{1}{2}(|H_1'\rangle \otimes |H_2'\rangle \otimes |V_3'\rangle + |H_1'\rangle \otimes |V_2'\rangle \otimes |H_3'\rangle \\ &\quad + |V_1'\rangle|H_2'\rangle|H_3'\rangle + |V_1'\rangle|V_2'\rangle|V_3'\rangle) \end{aligned}$$

where

$$\hat{\sigma}_x \otimes \hat{\sigma}_x \otimes \hat{\sigma}_x |\Psi_{GHZ}^{xxx(-)}\rangle = -|\Psi_{GHZ}^{xxx(-)}\rangle.$$

Similarly, in quantum mechanics we get

$$\hat{\sigma}_{1x} \otimes \hat{\sigma}_{2x} \otimes \hat{\sigma}_{3x} |H_1'\rangle \otimes |H_2'\rangle \otimes |H_3'\rangle = |H_1'\rangle \otimes |H_2'\rangle \otimes |H_3'\rangle$$

$$\hat{\sigma}_{1x} \otimes \hat{\sigma}_{2x} \otimes \hat{\sigma}_{3x} |V_1'\rangle \otimes |V_2'\rangle \otimes |H_3'\rangle = |V_1'\rangle \otimes |V_2'\rangle \otimes |H_3'\rangle$$

$$\hat{\sigma}_{1x} \otimes \hat{\sigma}_{2x} \otimes \hat{\sigma}_{3x} |V_1'\rangle \otimes |H_2'\rangle \otimes |V_3'\rangle = |V_1'\rangle \otimes |H_2'\rangle \otimes |V_3'\rangle$$

$$\hat{\sigma}_{1x} \otimes \hat{\sigma}_{2x} \otimes \hat{\sigma}_{3x} |H_1'\rangle \otimes |V_2'\rangle \otimes |V_3'\rangle = |H_1'\rangle \otimes |V_2'\rangle \otimes |V_3'\rangle$$

Then the four states

$$|H_1'\rangle \otimes |H_2'\rangle \otimes |H_3'\rangle, |V_1'\rangle \otimes |V_2'\rangle \otimes |H_3'\rangle,$$

$$|V_1'\rangle \otimes |H_2'\rangle \otimes |V_3'\rangle, |H_1'\rangle \otimes |V_2'\rangle \otimes |V_3'\rangle$$

have the same eigenvalue (+1).

Thus any superposition of these states has the eigenvalue (+1). However, there is only one GHZ state, among these states, which is defined by

$$\begin{aligned} |\Psi_{GHZ}^{xxx(+)}\rangle &= \frac{1}{\sqrt{2}}[|x_1\rangle|x_2\rangle|x_3\rangle + |y_1\rangle|y_2\rangle|y_3\rangle] \\ &= \frac{1}{2}(|H_1'\rangle|H_2'\rangle|H_3'\rangle + |H_1'\rangle|V_2'\rangle|V_3'\rangle + |V_1'\rangle|H_2'\rangle|V_3'\rangle + |V_1'\rangle|V_2'\rangle|H_3'\rangle) \end{aligned}$$

where


$$\hat{\sigma}_x \otimes \hat{\sigma}_x \otimes \hat{\sigma}_x |\Psi_{GHZ}^{xxx(+)}\rangle = + |\Psi_{GHZ}^{xxx(+)}\rangle.$$

In quantum mechanics,  $|\Psi_{GHZ}^{xxx(+)}\rangle$  is the eigenket of  $\hat{\sigma}_{1x} \otimes \hat{\sigma}_{2x} \otimes \hat{\sigma}_{3x}$  with the eigenvalue of +1, since

$$\hat{\sigma}_{1x} \otimes \hat{\sigma}_{2x} \otimes \hat{\sigma}_{3x} |\Psi_{GHZ}^{xxx(+)}\rangle = |\Psi_{GHZ}^{xxx(+)}\rangle$$

where

$$\begin{aligned} |\Psi_{GHZ}^{xxx(+)}\rangle &= \frac{1}{\sqrt{2}}[|x_1\rangle|x_2\rangle|x_3\rangle + |y_1\rangle|y_2\rangle|y_3\rangle] \\ &= \frac{1}{2}(|H_1'\rangle|H_2'\rangle|H_3'\rangle + |H_1'\rangle|V_2'\rangle|V_3'\rangle + |V_1'\rangle|H_2'\rangle|V_3'\rangle + |V_1'\rangle|V_2'\rangle|H_3'\rangle) \end{aligned}$$




**Fig.** Fraction. Prediction from **the quantum mechanics**. There are four states denoted by  $|H_1'\rangle|H_2'\rangle|H_3'\rangle$ ,  $|H_1'\rangle|V_2'\rangle|V_3'\rangle$ ,  $|V_1'\rangle|H_2'\rangle|V_3'\rangle$ , and  $|V_1'\rangle|V_2'\rangle|H_3'\rangle$ , which are different from the states predicted from the local hidden theory. (**Pan et.al.**)

### 11. Detail of the GHZ experiment

We consider the GHZ state given by


$$\begin{aligned} |\psi_{GHZ}^{yx(+)}\rangle = & \frac{1}{2} [ |R_1\rangle \otimes |L_2\rangle \otimes |H_3'\rangle + |L_1\rangle \otimes |R_2\rangle \otimes |H_3'\rangle \\ & + |R_1\rangle \otimes |R_2\rangle \otimes |V_3'\rangle + |L_1\rangle \otimes |L_2\rangle \otimes |V_3'\rangle ] \end{aligned}$$

For polarization measurements in the  $L/R$  basis, the photons in modes 1 and 2 have equal probability for the combinations  $|R_1\rangle \otimes |L_2\rangle$ ,  $|L_1\rangle \otimes |R_2\rangle$ ,  $|R_1\rangle \otimes |R_2\rangle$ , and  $|L_1\rangle \otimes |L_2\rangle$ . If  $|R_1\rangle \otimes |R_2\rangle$  is obtained, the photon in mode 3 has to be in the state  $|V_3'\rangle$ .



**Fig.** Set up for the creation of a GHZ state using two pairs of polarization entangled photon (**Pan et al.**). BS: half beam splitter. PBS: polarized beam splitter. POL: polarizer.  $\lambda/4$  quarter wave plate. F: narrow bandwidth filter.  $\lambda/2$ : half wave plate. The half wave plate switches  $|y\rangle$  to  $|H'\rangle = \frac{1}{\sqrt{2}}[|x\rangle + |y\rangle]$ . Quarter wave plates and polarizer just before the detectors are used for correlation measurements.

This figure shows the experimental results for this correlation measurement. Quarter wave plates and polarizers just before detectors  $D_1$ ,  $D_2$ , and  $D_3$  in Fig. are set to  $|R_1\rangle \otimes |R_2\rangle \otimes |V_3'\rangle$  or  $|R_1\rangle \otimes |R_2\rangle \otimes |H_3'\rangle$ . The results clearly confirms the strong correlations of  $|R_1\rangle \otimes |R_2\rangle \otimes |V_3'\rangle$  in comparison to  $|R_1\rangle \otimes |R_2\rangle \otimes |H_3'\rangle$ .



**Fig.** A typical experimental result used in the GHZ argument. This is the  $yyx$  experiment measuring circular polarization on photons 1 and 2 and linear polarization on the third. (Pan et.al.)

For photon  $i$  we call these elements of reality  $X_i$  with values  $+1$  ( $-1$ ) for  $|H\rangle$  ( $|V\rangle$ ) polarizations and  $Y_i$  with values  $+1$  ( $-1$ ) for  $|R\rangle$  ( $|L\rangle$ ); we thus obtain the relations

(a) The relation:  $Y_1 Y_2 X_3 = -1$ ,

form

	Each eigenvalues			Resultant eigenvalue
$ R_1\rangle \otimes  L_2\rangle \otimes  H_3\rangle$	1	-1	1	-1
$ L_1\rangle \otimes  R_2\rangle \otimes  H_3\rangle$	-1	1	1	-1
$ R_1\rangle \otimes  R_2\rangle \otimes  V_3\rangle$	1	1	-1	-1

in order to be able to reproduce the quantum predictions of equation

$$|\psi_{GHZ}^{yx(+)}\rangle = \frac{1}{2} [ |R_1\rangle \otimes |L_2\rangle \otimes |H_3\rangle + |L_1\rangle \otimes |R_2\rangle \otimes |H_3\rangle + |R_1\rangle \otimes |R_2\rangle \otimes |V_3\rangle + |L_1\rangle \otimes |L_2\rangle \otimes |V_3\rangle ]$$

(b) The relation  $Y_1 X_2 Y_3 = -1$ ,

from

	Each eigenvalues			Resultant eigenvalue
$ L_1\rangle \otimes  H_2\rangle \otimes  R_3\rangle$	-1	1	1	-1 (eigenvalue)
$ R_1\rangle \otimes  H_2\rangle \otimes  L_3\rangle$	1	1	-1	-1
$ R_1\rangle \otimes  V_2\rangle \otimes  R_3\rangle$	1	-1	1	-1
$ L_1\rangle \otimes  V_2\rangle \otimes  L_3\rangle$	-1	-1	-1	-1

in order to be able to reproduce the quantum predictions of equation

$$|\psi_{GHZ}^{xy(+)}\rangle = \frac{1}{2} [ |L_1\rangle \otimes |H_2\rangle \otimes |R_3\rangle + |R_1\rangle \otimes |H_2\rangle \otimes |L_3\rangle + |R_1\rangle \otimes |V_2\rangle \otimes |R_3\rangle + |L_1\rangle \otimes |V_2\rangle \otimes |L_3\rangle ]$$

(c) The relation  $X_1 Y_2 Y_3 = -1$ ,

from

	Each eigenvlaues			Resultant eigenvalue
$ R_1\rangle \otimes  L_2\rangle \otimes  H_3'\rangle$	1	-1	1	-1
$ L_1\rangle \otimes  R_2\rangle \otimes  H_3'\rangle$	-1	1	1	-1
$ R_1\rangle \otimes  R_2\rangle \otimes  V_3'\rangle$	1	1	-1	-1
$ L_1\rangle \otimes  L_2\rangle \otimes  V_3'\rangle$	-1	-1	-1	-1

in order to be able to reproduce the quantum predictions of equation

$$\begin{aligned} |\Psi_{GHZ}^{yyx(+)}\rangle = \frac{1}{2} [ & |R_1\rangle \otimes |L_2\rangle \otimes |H_3'\rangle + |L_1\rangle \otimes |R_2\rangle \otimes |H_3'\rangle \\ & + |R_1\rangle \otimes |R_2\rangle \otimes |V_3'\rangle + |L_1\rangle \otimes |L_2\rangle \otimes |V_3'\rangle ] \end{aligned}$$

Because of Einstein locality any specific measurement for  $X$  must be independent of whether an  $X$  or  $Y$  measurement is performed on the other photon. As  $Y_1Y_1 = Y_2Y_2 = Y_3Y_3 = 1$ , we can write

$$X_1X_2X_3 = (X_1Y_2Y_3)(Y_1X_2Y_3)(Y_1Y_2X_3) = (-1)(-1)(-1) = -1 \quad (1)$$

We now consider a fourth experiment measuring linear  $|H'\rangle/|V'\rangle$  polarization on all three photons, that is, an  $xxx$  experiment. We investigate the possible outcomes that will be predicted by local realism based on the elements of reality introduced to explain the  $xxx$  experiments. We obtain

$$X_1X_2X_3 = 1 \quad (2)$$

from

	Each eigenvlaue			Resultant eigenvalue
$ H_1'\rangle  H_2'\rangle  H_3'\rangle$	1	1	1	1
$ H_1'\rangle  V_2'\rangle  V_3'\rangle$	1	-1	-1	1
$ V_1'\rangle  H_2'\rangle  V_3'\rangle$	-1	1	-1	1
$ V_1'\rangle  V_2'\rangle  H_3'\rangle$	-1	-1	1	1

in order to be able to reproduce the quantum predictions of equation

$$|\Psi_{GHZ}^{xxx(+)}\rangle = \frac{1}{2} (|H_1'\rangle |H_2'\rangle |H_3'\rangle + |H_1'\rangle |V_2'\rangle |V_3'\rangle + |V_1'\rangle |H_2'\rangle |V_3'\rangle + |V_1'\rangle |V_2'\rangle |H_3'\rangle).$$



with

$$\hat{\sigma}_x \otimes \hat{\sigma}_x \otimes \hat{\sigma}_x \left| \Psi_{GHZ}^{xxx(+)} \right\rangle = + \left| \Psi_{GHZ}^{xxx(+)} \right\rangle.$$

Then the value  $X_1 X_2 X_3 = -1$  from Eq.(1) is incompatible with the value  $X_1 X_2 X_3 = 1$  from Eq.(2), predicted from the quantum mechanics.

## 12. Eigenstates of $\hat{\sigma}_x \otimes \hat{\sigma}_x \otimes \hat{\sigma}_x$

$$\hat{\sigma}_x \otimes \hat{\sigma}_x \otimes \hat{\sigma}_x =$$

$$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

There are eight eigenstates. The eigenvalues of the four states is -1 and the eigenvalue of the four states is (+1).

- (i) Four states are the eigenkets of  $\hat{\sigma}_x \otimes \hat{\sigma}_x \otimes \hat{\sigma}_x$  with the eigenvalue (-1). One of these states is the GHZ state

$$\hat{\sigma}_x \otimes \hat{\sigma}_x \otimes \hat{\sigma}_x \left| \Psi_{GHZ}^{xxx(-)} \right\rangle = - \left| \Psi_{GHZ}^{xxx(-)} \right\rangle$$

$$\begin{aligned} |\psi_1\rangle &= \left| \Psi_{GHZ}^{xxx(-)} \right\rangle \\ &= \frac{1}{\sqrt{2}} (|x_1\rangle|x_2\rangle|x_3\rangle - |y_1\rangle|y_2\rangle|y_3\rangle) \\ &= \frac{1}{2} (|H_1'\rangle|H_2'\rangle|V_3'\rangle + |H_1'\rangle|V_2'\rangle|H_3'\rangle + |V_1'\rangle|H_2'\rangle|H_3'\rangle + |V_1'\rangle|V_2'\rangle|V_3'\rangle) \end{aligned}$$

$$\begin{aligned}
|\psi_2\rangle &= \frac{1}{\sqrt{2}}(|x_1\rangle|x_2\rangle|y_3\rangle - |y_1\rangle|y_2\rangle|x_3\rangle) \\
&= \frac{1}{2}(-|H_1'\rangle|H_2'\rangle|V_3'\rangle + |H_1'\rangle|V_2'\rangle|H_3'\rangle + |V_1'\rangle|H_2'\rangle|H_3'\rangle - |V_1'\rangle|V_2'\rangle|V_3'\rangle)
\end{aligned}$$

$$\begin{aligned}
|\psi_3\rangle &= \frac{1}{\sqrt{2}}(|x_1\rangle|y_2\rangle|x_3\rangle - |y_1\rangle|x_2\rangle|y_3\rangle) \\
&= \frac{1}{2}(|H_1'\rangle|H_2'\rangle|V_3'\rangle - |H_1'\rangle|V_2'\rangle|H_3'\rangle + |V_1'\rangle|H_2'\rangle|H_3'\rangle - |V_1'\rangle|V_2'\rangle|V_3'\rangle)
\end{aligned}$$

$$\begin{aligned}
|\psi_4\rangle &= \frac{1}{\sqrt{2}}(|x_1\rangle|y_2\rangle|y_3\rangle - |y_1\rangle|x_2\rangle|x_3\rangle) \\
&= \frac{1}{2}(-|H_1'\rangle|H_2'\rangle|V_3'\rangle - |H_1'\rangle|V_2'\rangle|H_3'\rangle + |V_1'\rangle|H_2'\rangle|H_3'\rangle + |V_1'\rangle|V_2'\rangle|V_3'\rangle)
\end{aligned}$$

- (ii) Four states are the eigenkets of  $\hat{\sigma}_x \otimes \hat{\sigma}_x \otimes \hat{\sigma}_x$  with the eigenvalue (+1). One of these states is the GHZ state

$$\hat{\sigma}_x \otimes \hat{\sigma}_x \otimes \hat{\sigma}_x |\Psi_{GHZ}^{xxx(+)}\rangle = + |\Psi_{GHZ}^{xxx(+)}\rangle$$

$$\begin{aligned}
|\psi_5\rangle &= |\Psi_{GHZ}^{xxx(+)}\rangle \\
&= \frac{1}{\sqrt{2}}(|x_1\rangle|x_2\rangle|x_3\rangle + |y_1\rangle|y_2\rangle|y_3\rangle) \\
&= \frac{1}{2}(|H_1'\rangle|H_2'\rangle|H_3'\rangle + |H_1'\rangle|V_2'\rangle|V_3'\rangle + |V_1'\rangle|H_2'\rangle|V_3'\rangle + |V_1'\rangle|V_2'\rangle|H_3'\rangle)
\end{aligned}$$

$$\begin{aligned}
|\psi_6\rangle &= \frac{1}{\sqrt{2}}(|x_1\rangle|x_2\rangle|y_3\rangle + |y_1\rangle|y_2\rangle|x_3\rangle) \\
&= \frac{1}{2}(|H_1'\rangle|H_2'\rangle|H_3'\rangle - |H_1'\rangle|V_2'\rangle|V_3'\rangle - |V_1'\rangle|H_2'\rangle|V_3'\rangle + |V_1'\rangle|V_2'\rangle|H_3'\rangle)
\end{aligned}$$

$$\begin{aligned}
|\psi_7\rangle &= \frac{1}{\sqrt{2}}(|x_1\rangle|y_2\rangle|x_3\rangle + |y_1\rangle|x_2\rangle|y_3\rangle) \\
&= \frac{1}{2}(|H_1'\rangle|H_2'\rangle|H_3'\rangle - |H_1'\rangle|V_2'\rangle|V_3'\rangle + |V_1'\rangle|H_2'\rangle|V_3'\rangle - |V_1'\rangle|V_2'\rangle|H_3'\rangle)
\end{aligned}$$

$$\begin{aligned}
|\psi_8\rangle &= \frac{1}{\sqrt{2}}(|x_1\rangle|y_2\rangle|y_3\rangle + |y_1\rangle|x_2\rangle|x_3\rangle) \\
&= \frac{1}{2}(|H_1'\rangle|H_2'\rangle|H_3'\rangle + |H_1'\rangle|V_2'\rangle|V_3'\rangle - |V_1'\rangle|H_2'\rangle|V_3'\rangle - |V_1'\rangle|V_2'\rangle|H_3'\rangle)
\end{aligned}$$

---

### 13. Eigenvalue problem $\hat{\sigma}_x \otimes \hat{\sigma}_y \otimes \hat{\sigma}_y$

$$\hat{\sigma}_x \otimes \hat{\sigma}_y \otimes \hat{\sigma}_y =$$

$$\begin{pmatrix}
0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 \\
0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\
0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\
0 & 0 & 0 & 0 & -1 & 0 & 0 & 0 \\
0 & 0 & 0 & -1 & 0 & 0 & 0 & 0 \\
0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\
0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\
-1 & 0 & 0 & 0 & 0 & 0 & 0 & 0
\end{pmatrix}$$

There are eight eigenstates. The eigenvalues of the four states is -1 and the eigenvalue of the four states is (+1).

- (i) Four states are the eigenkets of  $\hat{\sigma}_x \otimes \hat{\sigma}_y \otimes \hat{\sigma}_y$  with the eigenvalue (-1). One of these states is the GHZ state  $|\Psi_{GHZ}^{xxx(+)}\rangle$ ,

$$\begin{aligned}
|\psi_1\rangle &= |\Psi_{GHZ}^{xxx(+)}\rangle \\
&= \frac{1}{\sqrt{2}}(|x_1\rangle|x_2\rangle|x_3\rangle + |y_1\rangle|y_2\rangle|y_3\rangle) \\
&= \frac{1}{2}(|H_1'\rangle|L_2\rangle|R_3\rangle + |H_1'\rangle|R_2\rangle|L_3\rangle + |V_1'\rangle|L_2\rangle|L_3\rangle + |V_1'\rangle|R_2\rangle|R_3\rangle)
\end{aligned}$$

$$\begin{aligned}
|\psi_2\rangle &= \frac{i}{\sqrt{2}}(|x_1\rangle|x_2\rangle|y_3\rangle - |y_1\rangle|y_2\rangle|x_3\rangle) \\
&= \frac{1}{2}(|H_1'\rangle|L_2\rangle|R_3\rangle - |H_1'\rangle|R_2\rangle|L_3\rangle - |V_1'\rangle|L_2\rangle|L_3\rangle + |V_1'\rangle|R_2\rangle|R_3\rangle)
\end{aligned}$$

$$\begin{aligned}
|\psi_3\rangle &= \frac{i}{\sqrt{2}}(|x_1\rangle|y_2\rangle|x_3\rangle - |y_1\rangle|x_2\rangle|y_3\rangle) \\
&= \frac{1}{2}(-|H_1'\rangle|L_2\rangle|R_3\rangle + |H_1'\rangle|R_2\rangle|L_3\rangle - |V_1'\rangle|L_2\rangle|L_3\rangle + |V_1'\rangle|R_2\rangle|R_3\rangle)
\end{aligned}$$

$$\begin{aligned}
|\psi_4\rangle &= -\frac{1}{\sqrt{2}}(|x_1\rangle|y_2\rangle|y_3\rangle + |y_1\rangle|x_2\rangle|x_3\rangle) \\
&= \frac{1}{2}(-|H_1'\rangle|L_2\rangle|R_3\rangle - |H_1'\rangle|R_2\rangle|L_3\rangle + |V_1'\rangle|L_2\rangle|L_3\rangle + |V_1'\rangle|R_2\rangle|R_3\rangle)
\end{aligned}$$

- (ii) The four states are the eigenkets of  $\hat{\sigma}_x \otimes \hat{\sigma}_y \otimes \hat{\sigma}_y$  with the eigenvalue (+1). One of these states is the GHZ state denoted by  $|\Psi_{GHZ}^{xxx(-)}\rangle$ ,

$$\begin{aligned}
|\psi_5\rangle &= |\Psi_{GHZ}^{xxx(-)}\rangle \\
&= \frac{1}{\sqrt{2}}(|x_1\rangle|x_2\rangle|x_3\rangle - |y_1\rangle|y_2\rangle|y_3\rangle) \\
&= \frac{1}{2}(|H_1'\rangle|L_2\rangle|L_3\rangle + |H_1'\rangle|R_2\rangle|R_3\rangle + |V_1'\rangle|L_2\rangle|R_3\rangle + |V_1'\rangle|R_2\rangle|L_3\rangle)
\end{aligned}$$

$$\begin{aligned}
|\psi_6\rangle &= \frac{i}{\sqrt{2}}(|x_1\rangle|x_2\rangle|y_3\rangle + |y_1\rangle|y_2\rangle|x_3\rangle) \\
&= \frac{1}{2}(-|H_1'\rangle|L_2\rangle|L_3\rangle + |H_1'\rangle|R_2\rangle|R_3\rangle + |V_1'\rangle|L_2\rangle|R_3\rangle - |V_1'\rangle|R_2\rangle|L_3\rangle)
\end{aligned}$$

$$\begin{aligned}
|\psi_7\rangle &= \frac{i}{\sqrt{2}}(|x_1\rangle|y_2\rangle|x_3\rangle + |y_1\rangle|x_2\rangle|y_3\rangle) \\
&= \frac{1}{2}(-|H_1'\rangle|L_2\rangle|L_3\rangle + |H_1'\rangle|R_2\rangle|R_3\rangle - |V_1'\rangle|L_2\rangle|R_3\rangle + |V_1'\rangle|R_2\rangle|L_3\rangle)
\end{aligned}$$

$$\begin{aligned}
|\psi_8\rangle &= \frac{1}{\sqrt{2}}(|x_1\rangle|y_2\rangle|y_3\rangle - |y_1\rangle|x_2\rangle|x_3\rangle) \\
&= \frac{1}{2}(-|H_1'\rangle|L_2\rangle|L_3\rangle - |H_1'\rangle|R_2\rangle|R_3\rangle + |V_1'\rangle|L_2\rangle|R_3\rangle + |V_1'\rangle|R_2\rangle|L_3\rangle)
\end{aligned}$$

#### 14. Eigenvalue problem $\hat{\sigma}_y \otimes \hat{\sigma}_x \otimes \hat{\sigma}_y$

$$\hat{\sigma}_y \otimes \hat{\sigma}_x \otimes \hat{\sigma}_y =$$

$$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

There are eight eigenstates. The eigenvalues of the four states is -1 and the eigenvalue of the four states is (+1).

- (i) Four states are the eigenkets of  $\hat{\sigma}_y \otimes \hat{\sigma}_x \otimes \hat{\sigma}_y$  with the eigenvalue (-1). One of these states is the GHZ state  $|\Psi_{GHZ}^{xxx(+)}\rangle$ ,

$$\begin{aligned} |\psi_1\rangle &= |\Psi_{GHZ}^{xxx(+)}\rangle \\ &= \frac{1}{\sqrt{2}}(|x_1\rangle|x_2\rangle|x_3\rangle + |y_1\rangle|y_2\rangle|y_3\rangle) \\ &= \frac{1}{2}(|H_1'\rangle|L_2\rangle|R_3\rangle + |H_1'\rangle|R_2\rangle|L_3\rangle + |V_1'\rangle|L_2\rangle|L_3\rangle + |V_1'\rangle|R_2\rangle|R_3\rangle) \end{aligned}$$

$$\begin{aligned} |\psi_2\rangle &= \frac{1}{\sqrt{2}}(|x_1\rangle|x_2\rangle|y_3\rangle - |y_1\rangle|y_2\rangle|x_3\rangle) \\ &= \frac{1}{2}(|H_1'\rangle|L_2\rangle|R_3\rangle - |H_1'\rangle|R_2\rangle|L_3\rangle - |V_1'\rangle|L_2\rangle|L_3\rangle + |V_1'\rangle|R_2\rangle|R_3\rangle) \end{aligned}$$

$$\begin{aligned} |\psi_3\rangle &= -\frac{1}{\sqrt{2}}(|x_1\rangle|y_2\rangle|x_3\rangle + |y_1\rangle|x_2\rangle|y_3\rangle) \\ &= \frac{1}{2}(|H_1'\rangle|L_2\rangle|L_3\rangle - |H_1'\rangle|R_2\rangle|R_3\rangle + |V_1'\rangle|L_2\rangle|R_3\rangle - |V_1'\rangle|R_2\rangle|L_3\rangle) \end{aligned}$$

$$\begin{aligned} |\psi_4\rangle &= \frac{1}{\sqrt{2}}(|x_1\rangle|y_2\rangle|y_3\rangle - |y_1\rangle|x_2\rangle|x_3\rangle) \\ &= \frac{1}{2}(-|H_1'\rangle|L_2\rangle|L_3\rangle - |H_1'\rangle|R_2\rangle|R_3\rangle + |V_1'\rangle|L_2\rangle|R_3\rangle + |V_1'\rangle|R_2\rangle|L_3\rangle) \end{aligned}$$

- (ii) The four states are the eigenkets of  $\hat{\sigma}_y \otimes \hat{\sigma}_x \otimes \hat{\sigma}_y$  with the eigenvalue (+1). One of these states is the GHZ state denoted by  $|\Psi_{GHZ}^{xxx(-)}\rangle$ ,

$$\begin{aligned} |\psi_5\rangle &= |\Psi_{GHZ}^{xxx(-)}\rangle \\ &= \frac{1}{\sqrt{2}}(|x_1\rangle|x_2\rangle|x_3\rangle - |y_1\rangle|y_2\rangle|y_3\rangle) \\ &= \frac{1}{2}(|H_1'\rangle|L_2\rangle|L_3\rangle + |H_1'\rangle|R_2\rangle|R_3\rangle + |V_1'\rangle|L_2\rangle|R_3\rangle + |V_1'\rangle|R_2\rangle|L_3\rangle) \end{aligned}$$

$$\begin{aligned} |\psi_6\rangle &= \frac{1}{\sqrt{2}}(|x_1\rangle|x_2\rangle|y_3\rangle + |y_1\rangle|y_2\rangle|x_3\rangle) \\ &= \frac{1}{2}(-|H_1'\rangle|L_2\rangle|L_3\rangle + |H_1'\rangle|R_2\rangle|R_3\rangle + |V_1'\rangle|L_2\rangle|R_3\rangle - |V_1'\rangle|R_2\rangle|L_3\rangle) \end{aligned}$$

$$\begin{aligned} |\psi_7\rangle &= \frac{1}{\sqrt{2}}(-|x_1\rangle|y_2\rangle|x_3\rangle + |y_1\rangle|x_2\rangle|y_3\rangle) \\ &= \frac{1}{2}(|H_1'\rangle|L_2\rangle|R_3\rangle - |H_1'\rangle|R_2\rangle|L_3\rangle + |V_1'\rangle|L_2\rangle|L_3\rangle - |V_1'\rangle|R_2\rangle|R_3\rangle) \end{aligned}$$

$$\begin{aligned} |\psi_8\rangle &= -\frac{1}{\sqrt{2}}(|x_1\rangle|y_2\rangle|y_3\rangle + |y_1\rangle|x_2\rangle|x_3\rangle) \\ &= \frac{1}{2}(-|H_1'\rangle|L_2\rangle|R_3\rangle - |H_1'\rangle|R_2\rangle|L_3\rangle + |V_1'\rangle|L_2\rangle|L_3\rangle + |V_1'\rangle|R_2\rangle|R_3\rangle) \end{aligned}$$

### 15. Eigenstates of $\hat{\sigma}_y \otimes \hat{\sigma}_y \otimes \hat{\sigma}_x$

$$\hat{\sigma}_y \otimes \hat{\sigma}_y \otimes \hat{\sigma}_x =$$

$$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

There are eight eigenstates. The eigenvalues of the four states is -1 and the eigenvalue of the four states is (+1).

- (i) Four states are the eigenkets of  $\hat{\sigma}_y \otimes \hat{\sigma}_y \otimes \hat{\sigma}_x$  with the eigenvalue (-1). One of these states is the GHZ state  $|\Psi_{GHZ}^{xxx(+)}\rangle$ , given by

$$\begin{aligned} |\psi_1\rangle &= |\Psi_{GHZ}^{xxx(+)}\rangle \\ &= \frac{1}{\sqrt{2}}(|x_1\rangle|x_2\rangle|x_3\rangle + |y_1\rangle|y_2\rangle|y_3\rangle) \\ &= \frac{1}{2}(|L_1\rangle|L_2\rangle|V_3'\rangle + |L_1\rangle|R_2\rangle|H_3'\rangle + |R_1\rangle|L_2\rangle|H_3'\rangle + |R_1\rangle|R_2\rangle|V_3'\rangle) \end{aligned}$$

The other three states are not the GHZ state;

$$\begin{aligned} |\psi_2\rangle &= -\frac{1}{\sqrt{2}}(|x_1\rangle|x_2\rangle|y_3\rangle + |y_1\rangle|y_2\rangle|x_3\rangle) \\ &= \frac{1}{2}(|L_1\rangle|L_2\rangle|V_3'\rangle - |L_1\rangle|R_2\rangle|H_3'\rangle - |R_1\rangle|L_2\rangle|H_3'\rangle + |R_1\rangle|R_2\rangle|V_3'\rangle) \end{aligned}$$

$$\begin{aligned} |\psi_3\rangle &= \frac{i}{\sqrt{2}}(|x_1\rangle|y_2\rangle|x_3\rangle + |y_1\rangle|x_2\rangle|y_3\rangle) \\ &= \frac{1}{2}(-|L_1\rangle|L_2\rangle|V_3'\rangle + |L_1\rangle|R_2\rangle|H_3'\rangle - |R_1\rangle|L_2\rangle|H_3'\rangle + |R_1\rangle|R_2\rangle|V_3'\rangle) \end{aligned}$$

$$\begin{aligned} |\psi_4\rangle &= \frac{i}{\sqrt{2}}(|x_1\rangle|y_2\rangle|y_3\rangle - |y_1\rangle|x_2\rangle|x_3\rangle) \\ &= \frac{1}{2}(|L_1\rangle|L_2\rangle|V_3'\rangle + |L_1\rangle|R_2\rangle|H_3'\rangle - |R_1\rangle|L_2\rangle|H_3'\rangle - |R_1\rangle|R_2\rangle|V_3'\rangle) \end{aligned}$$

- (ii) Four states are the eigenkets of  $\hat{\sigma}_y \otimes \hat{\sigma}_y \otimes \hat{\sigma}_x$  with the eigenvalue (+1). One of these state is the GHZ state  $|\Psi_{GHZ}^{xxx(-)}\rangle$ ,

$$\begin{aligned}
|\psi_5\rangle &= |\Psi_{GHZ}^{xxx(-)}\rangle \\
&= \frac{1}{\sqrt{2}}(|x_1\rangle|x_2\rangle|x_3\rangle - |y_1\rangle|y_2\rangle|y_3\rangle) \\
&= \frac{1}{2}(|L_1\rangle|L_2\rangle|H_3'\rangle + |L_1\rangle|R_2\rangle|V_3'\rangle + |R_1\rangle|L_2\rangle|V_3'\rangle + |R_1\rangle|R_2\rangle|H_3'\rangle)
\end{aligned}$$

$$\begin{aligned}
|\psi_6\rangle &= \frac{1}{\sqrt{2}}(-|x_1\rangle|x_2\rangle|y_3\rangle + |y_1\rangle|y_2\rangle|x_3\rangle) \\
&= \frac{1}{2}(-|L_1\rangle|L_2\rangle|H_3'\rangle + |L_1\rangle|R_2\rangle|V_3'\rangle + |R_1\rangle|L_2\rangle|V_3'\rangle - |R_1\rangle|R_2\rangle|H_3'\rangle)
\end{aligned}$$

$$\begin{aligned}
|\psi_7\rangle &= \frac{1}{\sqrt{2}}(i|x_1\rangle|y_2\rangle|x_3\rangle + |y_1\rangle|x_2\rangle|y_3\rangle) \\
&= \frac{1}{2}(-|L_1\rangle|L_2\rangle|H_3'\rangle + |L_1\rangle|R_2\rangle|V_3'\rangle - |R_1\rangle|L_2\rangle|V_3'\rangle + |R_1\rangle|R_2\rangle|H_3'\rangle)
\end{aligned}$$

$$\begin{aligned}
|\psi_8\rangle &= -i\frac{1}{\sqrt{2}}(|x_1\rangle|y_2\rangle|y_3\rangle + |y_1\rangle|x_2\rangle|x_3\rangle) \\
&= \frac{1}{2}(|L_1\rangle|L_2\rangle|H_3'\rangle + |L_1\rangle|R_2\rangle|V_3'\rangle - |R_1\rangle|L_2\rangle|V_3'\rangle - |R_1\rangle|R_2\rangle|H_3'\rangle)
\end{aligned}$$

## 16. Summary

(a) The GHZ state  $|\Psi_{GHZ}^{xxx(+)}\rangle$

$|\Psi_{GHZ}^{xxx(+)}\rangle$  is the simultaneous eigenket of

$$\hat{\sigma}_x \otimes \hat{\sigma}_y \otimes \hat{\sigma}_y |\Psi_{GHZ}^{xxx(+)}\rangle = -|\Psi_{GHZ}^{xxx(+)}\rangle,$$

$$\hat{\sigma}_y \otimes \hat{\sigma}_x \otimes \hat{\sigma}_y |\Psi_{GHZ}^{xxx(+)}\rangle = -|\Psi_{GHZ}^{xxx(+)}\rangle,$$

$$\hat{\sigma}_y \otimes \hat{\sigma}_y \otimes \hat{\sigma}_x |\Psi_{GHZ}^{xxx(+)}\rangle = -|\Psi_{GHZ}^{xxx(+)}\rangle,$$

$$\hat{\sigma}_x \otimes \hat{\sigma}_x \otimes \hat{\sigma}_x |\Psi_{GHZ}^{xxx(+)}\rangle = +|\Psi_{GHZ}^{xxx(+)}\rangle$$



(b) The GHZ state  $|\Psi_{GHZ}^{xxx(-)}\rangle$   
 $|\Psi_{GHZ}^{xxx(-)}\rangle$  is the simultaneous eigenket of

$$\hat{\sigma}_x \otimes \hat{\sigma}_y \otimes \hat{\sigma}_y |\Psi_{GHZ}^{xxx(-)}\rangle = + |\Psi_{GHZ}^{xxx(-)}\rangle$$

$$\hat{\sigma}_y \otimes \hat{\sigma}_x \otimes \hat{\sigma}_y |\Psi_{GHZ}^{xxx(-)}\rangle = + |\Psi_{GHZ}^{xxx(-)}\rangle$$

$$\hat{\sigma}_y \otimes \hat{\sigma}_y \otimes \hat{\sigma}_x |\Psi_{GHZ}^{xxx(-)}\rangle = + |\Psi_{GHZ}^{xxx(-)}\rangle$$

$$\hat{\sigma}_x \otimes \hat{\sigma}_x \otimes \hat{\sigma}_x |\Psi_{GHZ}^{xxx(-)}\rangle = - |\Psi_{GHZ}^{xxx(-)}\rangle$$

---

## 17. The GHS state in the configuration $yyx$ , $xyx$ , and $xyy$

(i)  $|\Psi_{GHZ}^{(+)}\rangle$  state

$$\begin{aligned} |\Psi_{GHZ}^{yyx(+)}\rangle &= \frac{1}{2} [ |R_1\rangle \otimes |L_2\rangle \otimes |H_3'\rangle + |L_1\rangle \otimes |R_2\rangle \otimes |H_3'\rangle \\ &\quad + |R_1\rangle \otimes |R_2\rangle \otimes |V_3'\rangle + |L_1\rangle \otimes |L_2\rangle \otimes |V_3'\rangle ] \end{aligned}$$

$$\begin{aligned} |\Psi_{GHZ}^{xyx(+)}\rangle &= \frac{1}{2} [ |L_1\rangle \otimes |H_2'\rangle \otimes |R_3\rangle + |R_1\rangle \otimes |H_2'\rangle \otimes |L_3\rangle \\ &\quad + |R_1\rangle \otimes |V_2'\rangle \otimes |R_3\rangle + |L_1\rangle \otimes |V_2'\rangle \otimes |L_3\rangle ] \end{aligned}$$

$$\begin{aligned} |\Psi_{GHZ}^{xyy(+)}\rangle &= \frac{1}{2} [ |H_1'\rangle \otimes |R_2\rangle \otimes |L_3\rangle + |H_1'\rangle \otimes |L_2\rangle \otimes |R_3\rangle \\ &\quad + |V_1'\rangle \otimes |R_2\rangle \otimes |R_3\rangle + |V_1'\rangle \otimes |L_2\rangle \otimes |L_3\rangle ] \end{aligned}$$

where

$$\hat{\sigma}_x \otimes \hat{\sigma}_y \otimes \hat{\sigma}_y |\Psi_{GHZ}^{xyy(+)}\rangle = - |\Psi_{GHZ}^{xyy(+)}\rangle.$$

$$\hat{\sigma}_y \otimes \hat{\sigma}_x \otimes \hat{\sigma}_y |\Psi_{GHZ}^{xyx(+)}\rangle = - |\Psi_{GHZ}^{xyx(+)}\rangle$$

$$\hat{\sigma}_y \otimes \hat{\sigma}_y \otimes \hat{\sigma}_x \left| \Psi_{GHZ}^{yyx(+)} \right\rangle = - \left| \Psi_{GHZ}^{yyx(+)} \right\rangle,$$

(ii)  $\left| \Psi_{GHZ}^{(-)} \right\rangle$  state

$$\left| \Psi_{GHZ}^{yyx(-)} \right\rangle = \frac{1}{2} (|L_1\rangle |L_2\rangle |H_3'\rangle + |R_1\rangle |R_2\rangle |H_3'\rangle + |L_1\rangle |R_2\rangle |V_3'\rangle + |R_1\rangle |L_2\rangle |V_3'\rangle)$$

$$\left| \Psi_{GHZ}^{xyx(-)} \right\rangle = \frac{1}{2} (|L_1\rangle |H_2'\rangle |L_3\rangle + |R_1\rangle |H_2'\rangle |R_3\rangle + |L_1\rangle |V_2'\rangle |R_3\rangle + |R_1\rangle |V_2'\rangle |L_3\rangle)$$

$$\left| \Psi_{GHZ}^{xyy(-)} \right\rangle = \frac{1}{2} (|H_1'\rangle |L_2\rangle |L_3\rangle + |H_1'\rangle |R_2\rangle |R_3\rangle + |V_1'\rangle |L_2\rangle |R_3\rangle + |V_1'\rangle |R_2\rangle |L_3\rangle)$$

where

$$\hat{\sigma}_x \otimes \hat{\sigma}_y \otimes \hat{\sigma}_y \left| \Psi_{GHZ}^{xyy(-)} \right\rangle = + \left| \Psi_{GHZ}^{xyy(-)} \right\rangle$$

$$\hat{\sigma}_y \otimes \hat{\sigma}_x \otimes \hat{\sigma}_y \left| \Psi_{GHZ}^{xyx(-)} \right\rangle = + \left| \Psi_{GHZ}^{xyx(-)} \right\rangle$$

$$\hat{\sigma}_y \otimes \hat{\sigma}_y \otimes \hat{\sigma}_x \left| \Psi_{GHZ}^{yyx(-)} \right\rangle = + \left| \Psi_{GHZ}^{yyx(-)} \right\rangle$$

---

## REFERENCES

- G. Greenstein and A.G. Zajon, *The Quantum Challenge*, 2<sup>nd</sup> edition (Jones and Bartlett, 2005).  
 J. Audretsch, *Entangled Systems, New Directions in Quantum Physics* (Wiley-VCH, 2007).  
 A. Peres, *Quantum Theory: Concepts and Methods* (Kluwer Academic Publishers, 2002).  
 Y. Aharonov and D. Rohrlich, *Quantum Paradoxes: Quantum Theory for the Perplexed* (Wiley-VCH, 2005).  
 A.D. Aczel, *Entanglement: The Greatest Mystery in Physics* (Four Walls Eight Windows, New York, 2002).  
 A. Furusawa and P. van Loock, *Quantum Teleportation and Entanglement* (Wiley-VCH, 2011).  
 S. Stenholm and K.-A. Suominen, *Quantum Approach to Information* (Wiley Interscience, 2005).  
 G. Jaeger, *Entanglement, Information, and the Interpretation of Quantum Mechanics* (Springer, 2009).  
 F. Laloë, *Do we understand Quantum Mechanics?* (Cambridge, 2012).

J.-W. Pan et al., Nature 403, p.515 (2000). Experimental test of quantum nonlocality in three-photon Greenberger-Horne-Zeilinger Entanglement.

G.J. Milburn, Feynman processor (Perseus Book, Cambridge MA, 1998).

R.A. Bertlmann and A. Zeilinger (eds.), Quantum [Un]speakable: From Bell to Quantum Information (Springer, 2002).

---

**APPENDIX-I. Eigenvalue problem  $\hat{\sigma}_x \otimes \hat{\sigma}_x \otimes \hat{\sigma}_y$**

$$\hat{\sigma}_x \otimes \hat{\sigma}_x \otimes \hat{\sigma}_y = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & -i \\ 0 & 0 & 0 & 0 & 0 & 0 & i & 0 \\ 0 & 0 & 0 & 0 & 0 & -i & 0 & 0 \\ 0 & 0 & 0 & 0 & i & 0 & 0 & 0 \\ 0 & 0 & 0 & -i & 0 & 0 & 0 & 0 \\ 0 & 0 & i & 0 & 0 & 0 & 0 & 0 \\ 0 & -i & 0 & 0 & 0 & 0 & 0 & 0 \\ i & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

There are eight eigenstates. The eigenvalues of the four states is -1 and the eigenvalue of the other four states is (+1). **There is no eigenket corresponding to the GHZ state.**

$$\begin{aligned} |\psi_1\rangle &= \frac{1}{\sqrt{2}}(i|x_1\rangle|x_2\rangle|x_3\rangle + |y_1\rangle|y_2\rangle|y_3\rangle) \\ &= \frac{1}{2}(i|H_1'\rangle|H_2'\rangle|R_3\rangle + i|H_1'\rangle|V_2'\rangle|L_3\rangle + i|V_1'\rangle|H_2'\rangle|L_3\rangle + i|V_1'\rangle|V_2'\rangle|R_3\rangle) \end{aligned}$$

$$\begin{aligned} |\psi_2\rangle &= \frac{1}{\sqrt{2}}(-|x_1\rangle|x_2\rangle|y_3\rangle + i|y_1\rangle|y_2\rangle|x_3\rangle) \\ &= \frac{1}{2}(-i|H_1'\rangle|H_2'\rangle|R_3\rangle + i|H_1'\rangle|V_2'\rangle|L_3\rangle + i|V_1'\rangle|H_2'\rangle|L_3\rangle - i|V_1'\rangle|V_2'\rangle|R_3\rangle) \end{aligned}$$

$$\begin{aligned} |\psi_3\rangle &= \frac{1}{\sqrt{2}}(|x_1\rangle|y_2\rangle|x_3\rangle - i|y_1\rangle|x_2\rangle|y_3\rangle) \\ &= \frac{1}{2}(|H_1'\rangle|H_2'\rangle|R_3\rangle - |H_1'\rangle|V_2'\rangle|L_3\rangle + |V_1'\rangle|H_2'\rangle|L_3\rangle - |V_1'\rangle|V_2'\rangle|R_3\rangle) \end{aligned}$$

$$\begin{aligned}
|\psi_4\rangle &= \frac{1}{\sqrt{2}}(i|x_1\rangle|y_2\rangle|y_3\rangle - |y_1\rangle|x_2\rangle|x_3\rangle) \\
&= \frac{1}{2}(-|H_1'\rangle|H_2'\rangle|R_3\rangle - |H_1'\rangle|V_2'\rangle|L_3\rangle + |V_1'\rangle|H_2'\rangle|L_3\rangle + |V_1'\rangle|V_2'\rangle|R_3\rangle)
\end{aligned}$$

(ii) The four states are the eigenkets of  $\hat{\sigma}_x \otimes \hat{\sigma}_x \otimes \hat{\sigma}_y$  with the eigenvalue (+1). There is no eigenkets corresponding to the GHZ state.

$$\begin{aligned}
|\psi_5\rangle &= \frac{1}{\sqrt{2}}(i|x_1\rangle|x_2\rangle|x_3\rangle - |y_1\rangle|y_2\rangle|y_3\rangle) \\
&= \frac{1}{2}(i|H_1'\rangle|H_2'\rangle|R_3\rangle + i|H_1'\rangle|V_2'\rangle|L_3\rangle + i|V_1'\rangle|H_2'\rangle|L_3\rangle + i|V_1'\rangle|V_2'\rangle|R_3\rangle)
\end{aligned}$$

$$\begin{aligned}
|\psi_6\rangle &= \frac{1}{\sqrt{2}}(-|x_1\rangle|x_2\rangle|y_3\rangle + i|y_1\rangle|y_2\rangle|x_3\rangle) \\
&= \frac{1}{2}(i|H_1'\rangle|H_2'\rangle|R_3\rangle - i|H_1'\rangle|V_2'\rangle|L_3\rangle - i|V_1'\rangle|H_2'\rangle|L_3\rangle + i|V_1'\rangle|V_2'\rangle|R_3\rangle)
\end{aligned}$$

$$\begin{aligned}
|\psi_7\rangle &= \frac{1}{\sqrt{2}}(|x_1\rangle|y_2\rangle|x_3\rangle + i|y_1\rangle|x_2\rangle|y_3\rangle) \\
&= \frac{1}{2}(|H_1'\rangle|H_2'\rangle|R_3\rangle - |H_1'\rangle|V_2'\rangle|L_3\rangle + |V_1'\rangle|H_2'\rangle|L_3\rangle - |V_1'\rangle|V_2'\rangle|R_3\rangle)
\end{aligned}$$

$$\begin{aligned}
|\psi_8\rangle &= -\frac{1}{\sqrt{2}}(i|x_1\rangle|y_2\rangle|y_3\rangle + |y_1\rangle|x_2\rangle|x_3\rangle) \\
&= \frac{1}{2}(-|H_1'\rangle|H_2'\rangle|R_3\rangle - |H_1'\rangle|V_2'\rangle|L_3\rangle + |V_1'\rangle|H_2'\rangle|L_3\rangle + |V_1'\rangle|V_2'\rangle|R_3\rangle)
\end{aligned}$$

## APPENDIX-II Eigenstates of $\hat{\sigma}_y \otimes \hat{\sigma}_y \otimes \hat{\sigma}_y$

$$\hat{\sigma}_y \otimes \hat{\sigma}_y \otimes \hat{\sigma}_y =$$

$$\begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 & 0 & i \\ 0 & 0 & 0 & 0 & 0 & 0 & -i & 0 \\ 0 & 0 & 0 & 0 & 0 & -i & 0 & 0 \\ 0 & 0 & 0 & 0 & i & 0 & 0 & 0 \\ 0 & 0 & 0 & -i & 0 & 0 & 0 & 0 \\ 0 & 0 & i & 0 & 0 & 0 & 0 & 0 \\ 0 & i & 0 & 0 & 0 & 0 & 0 & 0 \\ -i & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

There are eight eigenstates. The eigenvalues of the four states is -1 and the eigenvalue of the four states is (+1).

- (i) Four states are the eigenkets of  $\hat{\sigma}_y \otimes \hat{\sigma}_y \otimes \hat{\sigma}_y$  with the eigenvalue (-1). These states are not the GHZ state.

$$\begin{aligned} |\psi_1\rangle &= \frac{1}{\sqrt{2}}[|x_1\rangle|x_2\rangle|x_3\rangle + i|y_1\rangle|y_2\rangle|y_3\rangle] \\ &= \frac{1}{2}(|L_1\rangle|L_2\rangle|L_3\rangle + |L_1\rangle|R_2\rangle|R_3\rangle + |R_1\rangle|L_2\rangle|R_3\rangle + |R_1\rangle|R_2\rangle|L_3\rangle) \end{aligned}$$

$$\begin{aligned} |\psi_2\rangle &= \frac{1}{\sqrt{2}}[i|x_1\rangle|x_2\rangle|y_3\rangle + |y_1\rangle|y_2\rangle|x_3\rangle] \\ &= \frac{1}{2}(-|L_1\rangle|L_2\rangle|L_3\rangle + |L_1\rangle|R_2\rangle|R_3\rangle + |R_1\rangle|L_2\rangle|R_3\rangle - |R_1\rangle|R_2\rangle|L_3\rangle) \end{aligned}$$

$$\begin{aligned} |\psi_3\rangle &= \frac{1}{\sqrt{2}}[i|x_1\rangle|y_2\rangle|x_3\rangle + |y_1\rangle|x_2\rangle|y_3\rangle] \\ &= \frac{1}{2}(-|L_1\rangle|L_2\rangle|L_3\rangle + |L_1\rangle|R_2\rangle|R_3\rangle - |R_1\rangle|L_2\rangle|R_3\rangle + |R_1\rangle|R_2\rangle|L_3\rangle) \end{aligned}$$

$$\begin{aligned} |\psi_4\rangle &= \frac{1}{\sqrt{2}}[|x_1\rangle|y_2\rangle|y_3\rangle + i|y_1\rangle|x_2\rangle|x_3\rangle] \\ &= \frac{1}{2}(-|L_1\rangle|L_2\rangle|L_3\rangle - |L_1\rangle|R_2\rangle|R_3\rangle + |R_1\rangle|L_2\rangle|R_3\rangle + |R_1\rangle|R_2\rangle|L_3\rangle) \end{aligned}$$

- (ii) Four states are the eigenkets of  $\hat{\sigma}_y \otimes \hat{\sigma}_y \otimes \hat{\sigma}_y$  with the eigenvalue (+1). These states are not the GHZ state.

$$\begin{aligned} |\psi_5\rangle &= \frac{1}{\sqrt{2}}[|x_1\rangle|x_2\rangle|x_3\rangle - i|y_1\rangle|y_2\rangle|y_3\rangle] \\ &= \frac{1}{2}(|L_1\rangle|L_2\rangle|R_3\rangle + |L_1\rangle|R_2\rangle|L_3\rangle + |R_1\rangle|L_2\rangle|L_3\rangle + |R_1\rangle|R_2\rangle|R_3\rangle) \end{aligned}$$

$$\begin{aligned} |\psi_6\rangle &= \frac{1}{\sqrt{2}}[i|x_1\rangle|x_2\rangle|y_3\rangle - |y_1\rangle|y_2\rangle|x_3\rangle] \\ &= \frac{1}{2}(|L_1\rangle|L_2\rangle|R_3\rangle - |L_1\rangle|R_2\rangle|L_3\rangle - |R_1\rangle|L_2\rangle|L_3\rangle + |R_1\rangle|R_2\rangle|R_3\rangle) \end{aligned}$$

$$\begin{aligned} |\psi_7\rangle &= \frac{1}{\sqrt{2}}[i|x_1\rangle|y_2\rangle|x_3\rangle - |y_1\rangle|x_2\rangle|y_3\rangle] \\ &= \frac{1}{2}(-|L_1\rangle|L_2\rangle|R_3\rangle + |L_1\rangle|R_2\rangle|L_3\rangle - |R_1\rangle|L_2\rangle|L_3\rangle + |R_1\rangle|R_2\rangle|R_3\rangle) \end{aligned}$$

$$\begin{aligned} |\psi_7\rangle &= \frac{1}{\sqrt{2}}[|x_1\rangle|y_2\rangle|y_3\rangle - i|y_1\rangle|x_2\rangle|x_3\rangle] \\ &= \frac{1}{2}(|L_1\rangle|L_2\rangle|R_3\rangle + |L_1\rangle|R_2\rangle|L_3\rangle - |R_1\rangle|L_2\rangle|L_3\rangle - |R_1\rangle|R_2\rangle|R_3\rangle) \end{aligned}$$