

One dimensional bound state
Masatsugu Sei Suzuki
Department of Physics, SUNY at Binghamton
(Date: October 10, 2011)

1 One dimensional bound state

As a simple example of the calculation of discrete energy levels of a particle (with mass m) in quantum mechanics, we consider the one dimensional motion of a particle in the presence of a square-well potential barrier (width $2a$ and a depth V_0) as shown below.

$$V(x) = 0 \text{ for } |x| > a, \text{ and } -V_0 \text{ for } -a < x < a.$$

If the energy of the particle E is negative, the particle is confined and in a bound state. Here we discuss the energy eigenvalues and the eigenfunctions for the bound states from the solution of the Schrödinger equation.




Fig.8 One dimensional square well potential of width $2a$ and depth V_0 .

(a) The parity of the wave function

When potential is an even function (symmetric with respect to x), the wave function should have even parity or odd parity.

((Proof))

$$[\hat{\pi}, \hat{H}] = 0.$$

$\hat{\pi}$ is the parity operator.

$$\hat{\pi}^2 = 1 \quad \hat{\pi}^\dagger = \hat{\pi} = \hat{\pi}^{-1}.$$

$$\hat{\pi} \hat{x} \hat{\pi} = -\hat{x}. \quad \hat{\pi} \hat{p} \hat{\pi} = -\hat{p}.$$

\hat{H} is the Hamiltonian.

$$\hat{H} = \frac{\hat{p}^2}{2m} + V(\hat{x}),$$

and

$$\begin{aligned}\hat{\pi}\hat{H}\hat{\pi} &= \hat{\pi}\left[\frac{\hat{p}^2}{2m} + V(\hat{x})\right]\hat{\pi} \\ &= \frac{1}{2m}(\hat{\pi}\hat{p}\hat{\pi})^2 + V(\hat{\pi}\hat{x}\hat{\pi}) \\ &= \frac{1}{2m}(-\hat{p})^2 + V(-\hat{x}) \\ &= \frac{1}{2m}\hat{p}^2 + V(\hat{x})\end{aligned}$$

since $V(-\hat{x}) = V(\hat{x})$. Then we have a simultaneous eigenket:

$$\hat{H}|\psi\rangle = E|\psi\rangle, \text{ and } \hat{\pi}|\psi\rangle = \lambda|\psi\rangle.$$

Since $\hat{\pi}^2 = 1$,

$$\hat{\pi}^2|\psi\rangle = \lambda\hat{\pi}|\psi\rangle = \lambda^2|\psi\rangle = |\psi\rangle.$$

Thus we have $\lambda = \pm 1$.

or

$$\hat{\pi}|\psi\rangle = \pm|\psi\rangle,$$

$$\langle x|\hat{\pi}|\psi\rangle = \pm\langle x|\psi\rangle.$$

Since

$$\hat{\pi}|x\rangle = |-x\rangle, \text{ or } \langle x|\hat{\pi}^\dagger = \langle x|\hat{\pi} = \langle -x|$$

we have

$$\langle -x|\psi\rangle = \pm\langle x|\psi\rangle,$$

or

$$\psi(-x) = \pm\psi(x).$$

(b) Wavefunctions

In the Regions I, II, and III, the Schrödinger equation takes the form

$$\frac{d^2}{dx^2}\psi(x) - \kappa^2\psi(x) = 0 \quad \text{outside the well.}$$

$$\frac{d^2}{dx^2}\psi(x) + k^2\psi(x) = 0 \quad \text{inside the well.}$$

Here we define

$$\kappa^2 = \frac{2m}{\hbar^2}|E|, \quad k^2 = \frac{2m}{\hbar^2}(V_0 - |E|).$$

Here we introduce parameters (β and σ) for convenience,

$$\kappa^2 = \frac{2m}{\hbar^2}|E| = \frac{2mV_0}{\hbar^2} \frac{|E|}{V_0} = \frac{2mV_0a^2}{\hbar^2} \frac{1}{a^2} \frac{|E|}{V_0} = \frac{\beta^2}{a^2} \varepsilon,$$

or

$$\kappa^2 = \frac{\beta^2}{a^2} \varepsilon,$$

and

$$k^2 = \frac{2m}{\hbar^2}(V_0 - |E|) = \frac{2mV_0}{\hbar^2} \left(1 - \frac{|E|}{V_0}\right) = \frac{1}{a^2} \beta^2 (1 - \varepsilon),$$

where

$$\varepsilon = \frac{|E|}{V_0}, \quad \text{and} \quad \beta = \sqrt{\frac{2mV_0a^2}{\hbar^2}}.$$

We note that

$$k^2 + \kappa^2 = \frac{\beta^2}{a^2},$$

or

$$\xi^2 + \eta^2 = \beta^2,$$

where $ka = \xi$ and $ka = \eta$. The energy ε is given by

$$\varepsilon = \frac{\eta^2}{\beta^2} = 1 - \frac{\xi^2}{\beta^2}.$$

The stationary solution of the three regions are given by

$$\varphi_I(x) = Ae^{\kappa x},$$

$$\varphi_{II}(x) = B_1 e^{ikx} + B_2 e^{-ikx},$$

$$\varphi_{III}(x) = Ce^{-\kappa x}.$$

(i) The wave function with even parity

$$A = C,$$

$$B_1 = B_2 \equiv \frac{B}{2}.$$

The wavefunctions can be described by

$$\varphi_I(x) = Ae^{\kappa x},$$

$$\varphi_{II}(x) = B \cos(kx),$$

$$\varphi_{III}(x) = Ae^{-\kappa x}.$$

The derivatives are obtained by

$$\frac{d\varphi_I(x)}{dx} = A\kappa e^{\kappa x},$$

$$\frac{d\varphi_{II}(x)}{dx} = -Bk \sin(kx),$$

$$\frac{d\varphi_{III}(x)}{dx} = -A\kappa e^{-\kappa x}.$$

At $x = a$, $\varphi(x)$ and $\frac{d\varphi(x)}{dx}$ are continuous. Then we have

$$Ae^{-\kappa a} - B \cos(ka) = 0,$$

$$-A\kappa e^{-\kappa a} + Bk \sin(ka) = 0,$$

or

$$MX=0,$$

where

$$M = \begin{pmatrix} e^{-\kappa a} & -\cos(ka) \\ -\kappa e^{-\kappa a} & k \sin(ka) \end{pmatrix}, \quad X = \begin{pmatrix} A \\ B \end{pmatrix}.$$

The condition $\det M=0$ leads to

$$k \sin(ka)e^{-\kappa a} = \kappa e^{-\kappa a} \cos(ka),$$

or

$$\tan(ka) = \frac{\kappa}{k} \text{ for the even parity,}$$

or

$$\kappa a = ka \tan(ka) \quad \text{for the even parity.}$$

or

$$\eta = \xi \tan \xi.$$

The constants A, B, and C are given by

$$A = C = B e^{\kappa a} \cos(ka).$$

The condition of the normalization leads to the value of B.

(ii) The wave function with odd parity

$$A = -C,$$

$$B_1 = -B_2 \equiv \frac{B}{2i}.$$

The wavefunctions are given by

$$\varphi_I(x) = -Ae^{\kappa x},$$

$$\varphi_{II}(x) = B \sin(kx),$$

$$\varphi_{III}(x) = Ae^{-\kappa x}.$$

The derivatives are obtained as

$$\frac{d\varphi_I(x)}{dx} = -A\kappa e^{\kappa x},$$

$$\frac{d\varphi_{II}(x)}{dx} = Bk \cos(kx),$$

$$\frac{d\varphi_{III}(x)}{dx} = -A\kappa e^{-\kappa x}.$$

At $x = a$, $\varphi(x)$ and $\frac{d\varphi(x)}{dx}$ are continuous. Then we have

$$-Ae^{-\kappa a} + B \sin(ka) = 0,$$

$$-A\kappa e^{-\kappa a} - Bk \cos(ka) = 0,$$

or

$$MX=0,$$

where

$$M = \begin{pmatrix} -e^{-\kappa a} & \sin(ka) \\ -\kappa e^{-\kappa a} & -k \cos(\frac{ka}{2}) \end{pmatrix}, \quad X = \begin{pmatrix} A \\ B \end{pmatrix}.$$

The condition $\det M=0$ leads to

$$k \cos(ka)e^{-\kappa a} = -\kappa e^{-\kappa a} \sin(ka),$$

or

$$\kappa a = -ka \cot(ka) \quad \text{for the odd parity,}$$

or

$$\eta = -\xi \cot \xi .$$

We solve this eigenvalue problem using the Mathematica. The result is as follows.




Fig.9 Graphical solution. One solution with even parity for $0 < \beta < \pi/2$. One solution with even parity and one solution with odd parity for $\pi/2 < \beta < \pi$. Two solutions with even parity and one solution with odd parity for $\pi < \beta < 3\pi/2$. Two solutions with even parity and two solutions with odd parity for $3\pi/2 < \beta < 2\pi$. $\eta = \xi \tan \xi$ for the even parity (red lines). $\eta = -\xi \cot \xi$ for the odd parity (blue lines). The circles are denoted by $\xi^2 + \eta^2 = \beta^2$. The parameter β is changed as $\beta = 1, 2, 3, 4,$ and 5 . $\varepsilon = \frac{|E|}{V_0} = \frac{\eta^2}{\beta^2} = 1 - \frac{\xi^2}{\beta^2}$. $\xi = ka$ and $\eta = \kappa a$.

The normalized wavefunction for the even parity and odd parity are given by

$$\psi_{eI} = \frac{e^{\eta+x\eta} \cos[\xi]}{\sqrt{1 + \frac{\cos[\xi]^2}{\eta} + \frac{\sin[2\xi]}{2\xi}}}; \quad \psi_{eII} = \frac{\cos[x\xi]}{\sqrt{1 + \frac{\cos[\xi]^2}{\eta} + \frac{\sin[2\xi]}{2\xi}}};$$

$$\psi_{eIII} = \frac{e^{\eta-x\eta} \cos[\xi]}{\sqrt{1 + \frac{\cos[\xi]^2}{\eta} + \frac{\sin[2\xi]}{2\xi}}};$$

$$\psi_{oI} = -\frac{e^{\eta+x\eta} \sin[\xi]}{\sqrt{1 + \frac{\sin[\xi]^2}{\eta} - \frac{\sin[2\xi]}{2\xi}}}; \quad \psi_{oII} = \frac{\sin[x\xi]}{\sqrt{1 + \frac{\sin[\xi]^2}{\eta} - \frac{\sin[2\xi]}{2\xi}}};$$

$$\psi_{oIII} = \frac{e^{\eta-x\eta} \sin[\xi]}{\sqrt{1 + \frac{\sin[\xi]^2}{\eta} - \frac{\sin[2\xi]}{2\xi}}};$$

for the regions I, II, and III, where ψ_e is the wavefunction with the even parity and ψ_o is the wavefunction with the odd parity.

$\beta=1$

$\xi_{11} = 0.739085$	$\eta_{11} = 0.673612$	$\varepsilon_{11} = 0.453753$	even
-----------------------	------------------------	-------------------------------	------

$\beta=2$

$\xi_{21} = 1.02987$	$\eta_{21} = 1.71446$	$\varepsilon_{21} = 0.734844$	even
$\xi_{22} = 1.89549$	$\eta_{22} = 0.638045$	$\varepsilon_{22} = 0.101775$	odd

$\beta=3$

$\xi_{31} = 1.17012$	$\eta_{31} = 2.76239$	$\varepsilon_{31} = 0.847869$	even
$\xi_{32} = 2.27886$	$\eta_{32} = 1.9511$	$\varepsilon_{32} = 0.422976$	odd

$\beta=4$

$\xi_{41} = 1.25235$	$\eta_{41} = 3.7989$	$\varepsilon_{41} = 0.901976$	even
$\xi_{42} = 2.47458$	$\eta_{42} = 3.14269$	$\varepsilon_{42} = 0.617279$	odd
$\xi_{43} = 3.5953$	$\eta_{43} = 1.75322$	$\varepsilon_{43} = 0.192111$	even

$\beta = 5$

$\xi_{51} = 1.30644$	$\eta_{51} = 4.8263,$	$\varepsilon_{51} = 0.931729$	even
$\xi_{52} = 2.59574$	$\eta_{52} = 4.27342,$	$\varepsilon_{52} = 0.730486$	odd
$\xi_{53} = 3.83747$	$\eta_{53} = 3.20528,$	$\varepsilon_{53} = 0.410954$	even
$\xi_{54} = 4.9063$	$\eta_{54} = .963467,$	$\varepsilon_{54} = 0.0371307$	odd




Fig.10 Square well potential $V(x)$ of width $2a$ and depth V_0 . $\beta = 1$ and the corresponding wavefunction $\psi(x)$ which is normalized. There is one bound state (even parity) ($-\varepsilon_{11} = -0.45735$), where $\varepsilon = |E|/V_0$.




Fig.11 $\beta = 2$. There are two bound states. (i) The bound state (denoted by red) with even parity ($-\epsilon_{21} = -0.734844$). (ii) The bound state (denoted by blue) with odd parity ($-\epsilon_{22} = -0.101775$).




Fig.12 $\beta = 3$. There are two bound states. (i) The bound state (denoted by red) with even parity ($-\epsilon_{31} = -0.847869$). (ii) The bound state (denoted by blue) with odd parity ($-\epsilon_{32} = -0.422976$).




Fig.13 $\beta = 4$. There are three bound states. (i) The bound state (denoted by red) with even parity ($-\epsilon_{41} = -0.901976$). (ii) The bound state (denoted by blue)

with odd parity ($-\varepsilon_{42} = -0.617279$). (iii) The bound state (denoted by red) with even parity ($-\varepsilon_{43} = -0.192111$).





Fig.14 $\beta = 5$. There are four bound states. (i) The bound state (denoted by red) with even parity ($-\varepsilon_{51} = -0.931729$). (ii) The bound state (denoted by blue) with odd parity ($-\varepsilon_{52} = -0.730486$). (iii) The bound state (denoted by red) with even parity ($-\varepsilon_{53} = -0.410954$). (iv) The bound state (denoted by blue) with odd parity ($-\varepsilon_{54} = -0.0371307$).

REFERENCES

1. L.I. Schiff, Quantum Mechanics (McGraw-Hill, New York, 1955).
2. E. Merzbacher, Quantum Mechanics Third edition (John Wiley and Sons, New York, 1998).

II. Bound states

Bound state-II



We suppose that the potential energy is given by

$$V(x) = a|x| \quad (a > 0)$$

The Schrödinger equation is given by

$$\frac{d^2\psi}{dx^2} + \frac{2m}{\hbar^2}(E - a|x|)\psi = 0$$

where E is the energy of a particle with a mass m .

Since the potential is an even function of x , the wave function should be either an even function or an odd function of x .


The boundary condition for the wave function with odd parity;

$$\psi(x) = 0 \quad \text{at } x = 0.$$

The boundary condition for the wave function with even parity,

$$\frac{d\psi}{dx} = 0 \quad \text{at } x = 0.$$

$n = 0$:	even parity
$n = 1$	odd parity
$n = 2$	even parity
$n = 3$	odd parity
$n = 4$	even parity.



We use the dimensionless parameters;

$$\varepsilon = \frac{E}{(\hbar^2 a^2 / m)^{1/3}},$$

$$z = \frac{x}{(\hbar^2 / ma)^{1/3}}.$$

We note that

$$\frac{d\psi}{dx} = \frac{dz}{dx} \frac{d\psi}{dz} = \frac{1}{(\hbar^2 / ma)^{1/3}} \frac{d\psi}{dz}$$

$$\frac{d^2\psi}{dx^2} = \frac{dz}{dx} \frac{d}{dz} \frac{d\psi}{dx} = \frac{1}{(\hbar^2 / ma)^{2/3}} \frac{d^2\psi}{dz^2}$$

Then we get a differential equation

$$\frac{d^2\psi}{dz^2} + \frac{2m}{\hbar^2} (\hbar^2 / ma)^{2/3} [(\hbar^2 a^2 / m)^{1/3} \varepsilon - a|z|(\hbar^2 / ma)^{1/3}] \psi = 0$$

or

$$\frac{d^2\psi}{dz^2} + 2(\varepsilon - |z|)\psi = 0$$

Since the wave function either an even function or odd function, we consider the case of $z > 0$.

$$\frac{d^2\psi}{dz^2} + 2(\varepsilon - z)\psi = 0$$

Here we put $y = z - \varepsilon$.

$$\frac{d^2\psi(y)}{dy^2} - 2y\psi(y) = 0$$

((Mathematica))

```
Clear["Global`*"];
g1 = D[y[x], {x, 2}] - 2 x y[x] == 0;
eq1 = DSolve[g1, y[x], x]
{{y[x] -> AiryAi[2^(1/3) x] C[1] + AiryBi[2^(1/3) x] C[2]}}
```

`y1[x_] = y[x] /. eq1[[1]] /. {C[2] -> 0, C[1] -> 1}`
`AiryAi[2^(1/3) x]`

The solution of this differential equation is obtained as

$$\psi(y) = cA_i(2^{1/3} y)$$


where A_i is an Airy function and c is a constant to be determined from the normalization. Note that the second solution B_i is not a solution in this case since the function diverges when $x \rightarrow \infty$

(a) The case of even function (the even parity)

The boundary condition: $\frac{d\psi(z)}{dz}$ at $z = 0$.

which means that

$$\frac{d\psi(z)}{dz} = \frac{d\psi(y)}{dy} = 2^{1/3} A_i'(2^{1/3} y) = 0 \quad \text{at } y = -\varepsilon.$$



From this we get an energy eigenvalue for the wave functions with the even parity. The points with Green are located at $x = -\epsilon_0, -\epsilon_2, -\epsilon_4, -\epsilon_6, \dots$

$$\epsilon_0 = 0.808617$$

$$\psi_0(z) = 1.468 A_i(2^{1/3}(z - \epsilon_0))$$

$$\epsilon_2 = 2.5781$$

$$\psi_2(z) = 1.0510 A_i(2^{1/3}(z - \epsilon_2))$$

$$\epsilon_4 = 3.82572$$

$$\psi_4(z) = 1.0510 A_i(2^{1/3}(z - \epsilon_4))$$

$$\epsilon_6 = 4.89182$$

$$\epsilon_8 = 5.8513$$

$$\epsilon_{10} = 6.73732$$


$$\epsilon_{12} = 7.56829$$

$$\epsilon_{14} = 8.35581$$

(i)

$$\varepsilon_0 = 0.808617 \quad (n = 0);$$

even parity




The wave function is normalized.

(ii)

$$\varepsilon_2 = 2.5781 \quad (n = 2)$$

even parity




(b) The case of odd function:

The boundary condition: $\psi(z)$ at $z = 0$.

which means that

$$\psi(y) = 0 \quad \text{at } y = -\varepsilon.$$

From this we get an energy eigenvalue for the wave functions with the odd parity. The points with Green are located at $x = -\varepsilon_1, -\varepsilon_3, -\varepsilon_5, -\varepsilon_7, \dots$



$$\varepsilon_1 = 1.85576$$

$$\psi_1(z) = 1.1319 A_i(2^{1/3}(z - \varepsilon_1))$$

$$\varepsilon_3 = 3.24461$$

$$\psi_3(z) = 0.988282 A_i(2^{1/3}(z - \varepsilon_3))$$

$$\varepsilon_5 = 4.38167$$

$$\varepsilon_7 = 5.38661$$

$$\varepsilon_9 = 6.30526$$

$$\varepsilon_{11} = 7.16128$$


$$\varepsilon_{13} = 7.96889$$

(i)

$$\varepsilon_1 = 1.85576$$


$$(n = 1)$$

odd parity




(ii)

$\epsilon_3 = 3.24461$ $(n = 3)$ even parity



In summary we have



The plot of wave functions ($n = 0, 1, 2, 3$). The blue lines show the energy levels.